




ENCORE
 POUR-ON LOUSICIDE FOR SHEEP

DON'T

LET LICE

SPOIL

THE SHOW



Active Constituent: 35 g/L Imidacloprid
 Also contains: 300 g/L N-METHYL-2-PYRROLIDONE (solvent)

Claims: For the control of neonicotinoid susceptible body lice *Bovicola ovis* on shorn sheep up to 7 days off shears and unshorn lambs up to 2 months of age. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.

Container size: 20L

Applicator: Apply product only with the 6-stripe applicator.

Withholding Periods: MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: DO NOT USE on female sheep which are producing, or may in the future produce, milk or milk products for human consumption.

WOOL: DO NOT USE less than 6 months before shearing or fibre collection.

(ESI): DO NOT slaughter for export less than 63 days after treatment.


Other information on product:

May be used in conjunction with worm control drenches and at the same time as vaccination.

Contains a blue dye to identify treated sheep. This dye is scourable from the wool.

Is safe for use on breeding rams.

DOSAGE TABLE

BODYWEIGHT (KG)	DOSE (ML)	METHOD	APPLY AS BELOW
Below 6kg	8	Single Stripe	
6 – 8	12		
8.1 – 10	15		
10.1 – 12.5	20		
12.6 – 15	25		
15.1 – 20	30		
20.1 – 30	40	Double Stripe	
30.1 – 55	60		
55.1 – 80	80		
80.1 – 100	90		
Add 10mL for each 10kg above 100kg			