



io

independents Own

FLYK

SPRAY-ON SHEEP BLOWFLY TREATMENT

20L
AVAILABLE

Active:
50 g/L DICYCLANIL

NEW

BLOW

FLY STRIKE

AWAY!

READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

io
independents Own



FLYK

SPRAY-ON SHEEP BLOWFLY TREATMENT

ACTIVE CONSTITUENT: 50 g/L Dicyclanil

For the protection of sheep, either off-shears or with any length wool,
against fly strike (*Lucilia cuprina*) for 18 to 24 weeks.

For the protection of mulesing and marking wounds on sheep
against fly strike (*Lucilia cuprina*) during the wound healing process.



Made for AIRR
34 To Duntroon Rd.
Shepparton, VIC 3630
P: 03 5821 0922
ioindependentsown.com.au

CONTENTS
20
LITRES

io
independents Own

PROVED
OWNED
& MADE
IN AUSTRALIA



For the protection of
sheep, either off-shears
or with any length
wool, against fly strike
(*Lucilia cuprina*) for
18 to 24 weeks.

For the protection of
mulesing and marking
wounds on sheep
against fly strike
(*Lucilia cuprina*) during
the wound
healing process.

GENERAL DIRECTIONS:

io Flyk Spray-On provides season long protection to sheep against body, breach and poll strike and may be used either off-shears or on sheep with any length wool. *io Flyk Spray-On* also protects mulesing and marking wounds on lambs against fly strike during the wound healing process. The dose volume is determined by the weight of sheep being treated. Apply *io Flyk Spray-On* according to the diagrams and tables in the Directions for Use. DO NOT DILUTE. Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by The Hunter River Company. *io Flyk Spray-On* may be applied to all breeds of sheep to obtain protection against fly strike.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.

WOOL HARVEST INTERVAL: DO NOT USE less than 3 months before shearing or fibre collection.

TRADE ADVICE – EXPORT
SLAUGHTER INTERVAL (ESI): DO NOT use less than 120 days before slaughter for export.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

FOR LABEL, SDS AND ALL TECHNICAL
ENQUIRIES, PLEASE GO TO:

independentsown.com.au

E: info@airr.com.au

P: (03) 5821 0922