



Starvec Injection will treat and control internal parasites including gastrointestinal roundworms, lungworms, eyeworms, sucking and biting lice, mites and cattle tick in cattle.

Active Constituent: 10mg/ml Ivermectin

Available in the following sizes: 500ML

Applicator: 20ml Injector (Drencher with conversion kit)

Dosing & Administration: Use the contents within 24 hours of first broaching of the vial. Discard the unused portion.

The recommended dose is 1 mL STARMEC Ivermectin Injection for Cattle per 50kg liveweight (equivalent to 200 µg ivermectin per kg liveweight) by subcutaneous injection. DO NOT administer by the intravenous or intramuscular route.

Liveweight (kg) Dose volume (mL) Treatments per 500mL

Up to 100	2	250
101 – 150	3	166
151 – 200	4	125
201 – 250	5	100
251 – 300	6	83
301 – 350	7	71
351 – 400	8	62
401 – 450	9	55
451 – 500	10	50
501 – 550	11	45
551 – 600	12	41
601 – 650	13	38



Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active constituent: 20 g/L SPINOSAD.

Recommended applicator: iO 30ml Applicator with 4 hole T Bar Kit (31845).

Dosage & Administration: Apply iO SPINOFF POUR-ON evenly along the backline from the poll to the base of the tail using a manual or powered applicator with an iO SPINOFF POUR-ON 4-hole t-bar nozzle. Halve the dose volume and apply in 2 bands, approximately 5 cm from the mid-point of the backline on each side. Applying the product to the flanks will improve the spread of iO SPINOFF POUR-ON.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label



[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 5 g/L Moxidectin

Application: 30ml Variable Applicator with Pour-On Nozzle or Pour-On Applicator 65ml.

Dosing & Administration: MOXISTAR CATTLE POUR-ON should be applied along the top of the back from withers to the base of the tail. The volume to use is 1 mL per 10 kg liveweight. This will administer the correct dose of 0.5 mg/kg liveweight.

MOXISTAR CATTLE POUR-ON can be used in lactating dairy cattle, but DO NOT RETREAT LACTATING CATTLE AT LESS THAN 21 DAY INTERVALS.

Liveweight (kg) Dose (mL)

31-40	4
41-50	5
51-60	6
61-70	7
71-80	8

Liveweight (kg) Dose (mL)

81-90	9
91-100	10
101-125	12.5
126-150	15
151-175	17.5
176-200	20
201-250	25
251-300	30
301-350	35
351-400	40
401-450	45
451-500	50
501-550	55

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

For the treatment and control of moxidectin sensitive gastrointestinal parasites (including benzimidazole and/or levamisole resistant strains), lungworm and itch mite of sheep. An aid in the control of selenium responsive conditions.

Moxistar LV is highly effective against gastrointestinal parasites including Barber's Pole worm, Black scour worm, Stomach hair worm, Small brown stomach worm, Small intestinal worm, Thin necked intestinal worm, Intestinal threadworm, Nodule worm, Large mouthed bowel worm, Large lungworm & itch mite.

Active Constituent: 2 g/L Moxidectin 1 g/L Selenium (as Sodium selenate)

Dosage and Administration: MOXISTAR LV Se is ready to use. Administer 1 mL per 10 kg liveweight using any standard drenching equipment to give a dose of 0.2 mg moxidectin/kg and 0.1mg selenium/kg liveweight. Check dose rates and equipment before drenching commences.

Withholding Periods:

- MEAT: DO NOT USE less than 7 days before slaughter for human consumption.
- MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

- EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export.

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Containing moxidectin in a long-acting formulation, it provides sustained control of internal parasites in sheep during periods of challenge.

Active Ingredient: 20mg/ml Moxidectin

Benefits

- Provides control of internal parasites in sheep
- Contains moxidectin, a highly potent member of the macrocyclic lactone (ML) family
- Effective against parasites susceptible to ML drenches, including gastrointestinal roundworms and large lungworm
- ≥91 days persistent activity against Barber's Pole worm and Small Brown Stomach worm
- Up to 49 days persistent activity against Black Scour worm
- Up to 4 months protection during severe Barber's Pole worm challenge
- Long duration of activity may reduce the need for frequent drenching
- Suitable for use in highly contaminated pastures
- Controls nasal bot (larval stages) and itch mite
- Simple subcutaneous injection for ease of application

- Wide margin of safety when used as directed

When administered at the recommended dose rate, Moxistar LA Sheep Injection provides:

- At least 91 days persistent activity against Barber's Pole worm (*Haemonchus contortus*)
- At least 91 days persistent activity against Small Brown Stomach worm (*Teladorsagia circumcincta*)
- Up to 49 days persistent activity against Black Scour worm (*Trichostrongylus* spp.)
- During periods of severe Barber's Pole worm challenge, protection may extend for up to 4 months when used as directed.
- Provides long-acting control of key internal parasites in sheep, including extended protection against barber's pole worm and small brown stomach worm.
- Delivers persistent activity against black scour worm and helps protect sheep under heavy parasite challenge.

Withholding Period/ESI

- Meat: 91 days
- Milk: Do not use in female sheep which are producing, or may in the future produce, milk or milk products for human consumption
- ESI: 133 days

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

For the control of Blowfly Strike for up to 14 weeks. Will protect long wool sheep against fly strike for eleven weeks when used as directed.

Active Constituent: Cyromazine 60g/L.

Controls & Prevents: Will protect long wool sheep against fly strike for eleven weeks when used as directed. Indicated for the control of Blowfly Strike (*Lucilia cuprina*), which includes organophosphate resistant strains in long wool sheep for up to eleven weeks when used as directed.

Dose Rates & Administration Instructions: Please refer to package leaflet for directions on use. Do not dilute. Do not mix with any other product. Venus Spray-On may be applied to all breeds of sheep, providing protection against fly strike.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure



[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: Cyromazine 500g/L

Dose Rates & Administration Instructions: Venus Liquid is a water-soluble suspension for jetting, dipping and dressing.

Quick Facts & Benefits: The active ingredient in Venus Liquid is cyromazine. Cyromazine causes retardation, malformation and death of young fly larvae. Venus Liquid has a favourable environmental profile, and is safe for producers and sheep when used according to the label. Venus Liquid does not affect wool quality.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label



[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Designed for use where resistance to single actives may be present, Tripletec LV supports effective parasite control in modern livestock systems. By combining three actives from different chemical groups, it helps address worm populations that may not be adequately controlled by single-active drenches alone.

The low-volume formulation supports simple, accurate dosing (1 mL per 10 kg), helping improve consistency across mobs and reduce the risk of under-dosing.

Tripletec LV also includes selenium and cobalt to assist with trace element supplementation in sheep and cattle grazing in areas where these minerals may be deficient.

Active Ingredients

Levamisole Hydrochloride 80 g/L, Oxfendazole 45.3 g/L, Cobalt 5 g/L (as disodium cobalt EDTA), Abamectin 2 g/L, Selenium 1 g/L (as sodium selenate)

Key Benefits

- Triple-active, 3-way mineralised combination drench
- Broad internal parasite control for sheep and cattle
- Suitable for use where resistance to single actives may be present
- Low-volume formulation for simple, accurate dosing
- Supports integrated parasite management programs
- Includes selenium and cobalt for use in deficient areas

Effective Against


- Sheep & Cattle – Gastrointestinal Roundworms, including:
 - Barber's Pole Worm
 - Black Scour Worm
 - Small Brown Stomach Worm
 - Cooperia spp.
 - Nematodirus spp.
- and other common internal parasites

Dosage

1ml/10kg body weight by single oral administration. Refer to the label for further information

Withholding Periods

Meat: DO NOT USE less than 21 days before slaughter for human consumption.



Milk (Cattle): DO NOT USE in lactating cows or within 28 days of calving where milk or milk products will be used for human consumption. If cows calve earlier than 20 days after treatment, milk may contain residues. This milk MUST NOT BE USED for human consumption, or supplied for processing for at least 28 days following treatment. Calves fed this milk should not be slaughtered for human consumption within 21 days.

Export Slaughter Interval (ESI)

Cattle: DO NOT USE less than 21 days before slaughter for export.
Sheep: DO NOT USE less than 28 days before slaughter for export.

Before using this product, confirm the current ESI from the distributor on (03) 5820 8400 or the APVMA website (apvma.gov.au/residues).

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

iO Tick Off is a ready-to-use pour-on formulation. iO Tick Off breaks the tick life cycle. When treated with iO Tick Off, immature ticks are unable to molt to the next stage of their life cycle, and no larvae will hatch from eggs laid by treated females.

Active Constituent:

25g/L Fluazuron

Solvents:

100 g/L 1-dodecyl-2-pyrrolidone

100 g/L 1-octyl-2-pyrrolidone

415 g/L 1-methyl-2-pyrrolidone

Controls & Prevents: For strategic control of Cattle Tick, *Boophilu microplus*, (including all known acaricide resistant strains) on Beef Cattle.

Dose Rates & Administration Instructions: Apply iO Tick Off in two bands, roughly 7cm wide, one on each side of the spine between the rump and the shoulders. Only 3 maximum treatments per tick season annually. Make sure you re-treat when you first see ticks but not before at least six weeks after the previous treatment. Breed Category

Cattle



Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

iO Tick Off Duo contains fluazuron and ivermectin to give the dual benefits of tick control (fluazuron) and broad spectrum internal/external parasite protection (ivermectin) in one easy to use pour-on. Use in strategic cattle tick programs when broad spectrum parasite control is required in tick infested geographies.

Active Constituent:

15 g/L Fluazuron, 5 g/L Ivermectin

Controls & Prevents: Cattle Tick (*Rhipicephalus microplus*), ivermectin-sensitive gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, sucking and biting lice, sarcoptic and chorioptic mites, and buffalo fly (*Haematobia irritans exigua*) on Beef Cattle.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure



[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 5 g/L Ivermectin.

Controls & Prevents: For the treatment and control of ivermectin sensitive:

Gastrointestinal nematodes:

Adult and immature (L4): Small Brown Stomach Worm (including inhibited stages) (*Ostertagia ostertagi*), Barber's Pole Worm (*Haemonchus placei*), Stomach Hair Worm (*Trichostrongylus axei*), Intestinal Hair Worm (*T. colubriformis*), Small Intestinal Worm (*Cooperia* spp.), Nodule Worm (*Oesophagostomum radiatum*); and Adult stages of *Cooperia oncophora*, *C. punctata*, *Oesophagostomum venulosum*, Whip Worm (*Trichuris* spp.), Intestinal Threadworm (*Strongyloides papillosis*); and Immature stages (L4) of Thin Necked Intestinal Worm (*Nematodirus* spp.).


Lungworms: (adult and immature) *Dictyocaulus viviparus*.

Eyeworm: *Thelazia* spp. (adult stages).

Sucking and Biting Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Bovicola bovis*, *Solenopotes capillatus*.

Mites: *Chorioptes bovis*, *Sarcoptes scabiei* var *bovis*.

Buffalo Flies: *Haematobia irritans exigua* g. Cattle Ticks: (*Boophilus microplus*) including organophosphate, synthetic pyrethroid- and amidine-resistant strains.



Administration Instructions: This product is formulated only for external application to cattle; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead.

Dose Rates The dose is 1mL for each 10kg of liveweight. Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers etc.).
When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid overdosing. Do not underdose. Cattle should be weighed with either scales or with a weighband.

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia


1

Active constituents: Ivermectin 10g/L & Clorsulon 100g/L

Mectec Plus is a ready to use broad-spectrum endectocide injection for cattle containing Ivermectin and Clorsulon for the treatment and control of Ivermectin and Clorsulon sensitive strains of internal and external parasites of cattle, including adult liver flukes.

The recommended dose rate is 1mL/50kg live weight by subcutaneous injection. Heavier animals (for example mature bulls) should receive an additional 1 mL for each 50kg over 650kg.

Liveweight (kg)	Dose (mL)	No. of animals treated
Up to 100	2	250
101 – 150	3	166
151 – 200	4	125
201 – 250	5	100
251 – 300	6	83



301 – 350	7	71
Liveweight (kg)	Dose (mL)	No. of animals treated
351 – 400	8	62
401 – 450	9	55
451 – 500	10	50
501- 550	11	45
551 – 600	12	41
601 – 650	13	38

Withholding periods:

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: Zero (0) days

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT slaughter for export less than 42 days after treatment.

Recommended Applicators: iO 15 mL Injector

Storage: Store below 30c (Room temperature) and protect from light in a closed, original container.

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure



[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 16g/L Ivermectin

Dose Rates & Administration Instructions: JETTING FOR BLOWFLY STRIKE: When applied as directed by hand jetting or by modified Harrington jetting race JET AWAY BLOWFLY AND LICE JETTING FLUID FOR SHEEP protects sheep against strike by blowflies (*Lucilia cuprina*, *L. sericata*, *Chrysomya rufifacies* and *Calliphora nociva*) (including OP resistant strains of *Lucilia cuprina*) for 12 weeks under low to moderate fly pressure and will control maggots in existing strikes in the treated areas.

HAND DRESSING OF STRUCK SHEEP: JET AWAY can be applied at the blowfly dilution rate for treatment of blowfly strike wounds. Sheep should be sheared/clipped around the strike wound before saturating the area.

HAND JETTING FOR LICE: When applied as directed by hand-jetting JET AWAY also treats biting lice (*Bovicola ovis*), including synthetic pyrethroid resistant strains in long woolled sheep until the next shearing. Testing has confirmed that JET AWAY mixes with waters of various hardness. Use either a jetting handpiece with multiple nozzles that can be combed through the fleece or a "Dutjet" jetting wand. The jetting handpiece should deliver a penetrating jet of fluid without producing a spray or mist. A modified Harrington jetting race can also be used.



Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 50 g/L Dicyclanil

Controls & Prevents: For the protection of mulesing and marking wounds on sheep against fly strike (*Lucilia cuprina*) during the wound healing process.

Dose Rates: The dose volume is determined by the weight of sheep being treated.

Administration Instructions

Apply iO Flyk Spray-On according to the diagrams and tables in the Directions for Use. DO NOT DILUTE. Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by The Hunter River Company. iO Flyk Spray-On may be applied to all breeds of sheep to obtain protection against fly strike.

Significant soiling with dags, mud, dust or urine may reduce efficacy. Where significant dags are present, a light crutching is indicated prior to treatment. Protection period may be reduced if sheep scour heavily following application.

For the Prevention of Body Strike (Diagram 1)

Sheep Weight (kg)	Total Volume / Sheep (mL)	Volume per Treatment Band (mL)	No. of Bands	No. of Sheep Treated per 20L
10–20	14	7	2	1428
21–30	16	8	2	1250
31–50	20	10	2	1000
>50	24	12	2	833

For the Prevention of Breech Strike (Diagram 1)

Sheep Weight (kg)	Total Volume / Sheep (mL)	Volume per Treatment Band (mL)	No. of Bands	No. of Sheep Treated per 20L
10–20	7	7	1	2857
21–30	8	8	1	2500
31–50	10	10	1	2000
>50	12	12	1	1666

For the Prevention of Poll Strike (Diagram 1)

Rams and sheep susceptible to poll strike should be treated. Apply 5mL of iO Fly to the areas requiring protection, around the poll and horns.

For the Prevention of Fly Strike After Mulesing (Diagram 2)


Apply two bands, one over each side of the mulesing wound and overlapping 25mm onto the surrounding wool.

Sheep Weight (kg)	Total Volume / Sheep (mL)	Volume per Treatment Band (mL)	No. of Bands	No. of Sheep Treated per 20L
Up to 30	8	4	2	2500
31-50	10	5	2	2000

For the Prevention of Fly Strike After Marking (Diagram 3)


- When a tail has been amputated, apply 4mL over the crotch, ensuring sufficient coverage of the open wound.
- When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied.
- In the case of ram lambs, a further 4mL can be sprayed over and around the scrotum area after the rubber ring has been applied.

Lambs Weight (<30 kg)	Total Volume / Lamb (mL)	Volume per Treatment Band (mL)	No. of Bands	No. of Lambs Treated per 20L
Ram lambs	8	4	2	2500
Ewe lambs	4	4	1	5000



Critical Comments

- Use only a spray-on applicator gun recommended by The Hunter River Company.
- Ensure the applicator gun is clean.
- Calibrate applicator gun using an accurate volume measure.
- Ensure all areas to be protected are treated.

- 
- When preventing body strike ensure bands overlap along the midline.
 - When protecting mulesing wounds from fly strike, treat the wound and surrounding 25mm of wool.
 - When protecting marking wounds from fly strike, treat the wound and surrounding wool.
 - This product is not recommended for treating existing strikes. Use a registered product as a dressing.
 - Heavy rain following application may diminish protection.
 - iO Fly Spray-On contains a scourable colouring agent for easy identification.
 - Withholding periods must be observed following application.

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)



Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 65 g/L Dicyclanil

Available size: 20L

Dose Rates: The dose volume is determined by the weight of sheep being treated.

Withholding Periods:

Meat: Do not use less than 14 days before slaughter for human consumption.

ESI: DO NOT USE less than 9 weeks before slaughter or export.

Milk: Do not use on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.



Wool: Do not use less than 3 months before shearing or fibre collection.

Storage: Store below 30°C (Room Temperature). Do not freeze. Protect from light. Store in original container tightly closed in a dry, cool place.

Applicator: iO 30ml Variable Applicator

Recycle containers via [drumMUSTER](#)

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 10mg/ml Ivermectin and 75mg/ml Praziquantel

iO Equiduo (Ivermectin and Praziquantel) has been formulated for administration by oral drench or stomach tube (nasogastric intubation). One low-volume dose is effective against important internal parasites, including tapeworms and the arterial stages of *Strongylus vulgaris*, and bots.

Controls & Prevents: Treats Tapeworms, Roundworms/Ascarids, Small Strongylus, Large Strongylus, Intestinal Threadworms, Large Mouth Stomach Worms, Pinworms, Lungworms, Bots & Hairworms.

Dose Rates & Administration Instructions: Each 50mL bottle contains sufficient ivermectin and praziquantel to treat 5 x 500kg horses. The recommended dose is 1mL iO Equiduo per 50kg liveweight (equivalent to 200mg ivermectin per kg liveweight and 1.5 mg praziquantel per kg liveweight). iO Equiduo can be given as an oral drench or by stomach tube (nasogastric intubation). Nasogastric intubation should only be done by a veterinarian or under veterinary supervision.

Breed Category

Cattle



Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: Active Constituent: 100 g/L FENBENDAZOLE


Dose Rates & Administration Instructions:

CATTLE: 7.5 mg/kg Fenbendazole or 3 mL per 40 kg bodyweight. Cattle in excess of 680 kg to be dosed at 3 mL per 40 kg bodyweight.

HORSES: 10 mg/kg Fenbendazole or 5 mL per 50 kg body weight. For the control of *Strongyloides westeri* in foals: 50 mg/kg or 5 mL/10 kg body weight. Independents Own Fenbender 100 Oral Drench may be administered to horses by nasal tube, drenching gun or dosage syringe.

Bodyweight	Dose	No. of Animals per 1L Pack
------------	------	----------------------------

Up to 80 kg	6 mL	167
-------------	------	-----



81 – 120 kg	9 mL	133
121 – 160 kg	12 mL	83
161 – 200 kg	15 mL	67
201 – 240 kg	18 mL	55
241 – 280 kg	21 mL	47
281 – 320 kg	24 mL	41
321 – 360 kg	27 mL	37
361 – 400 kg	30 mL	33
401 – 440 kg	33 mL	30
441 – 480 kg	36 mL	27
481 – 520 kg	39 mL	25
521 – 560 kg	42 mL	23
561 – 600 kg	45 mL	22

Recycle containers via [drumMUSTER](#)

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download



Missing Attachment
Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 50 mg/g Praziquantel & 4 mg/g Abamectin. 35 ml syringe treats 700kg horse. Wide safety margin. Easy to administer with an Oatmeal Flavour.

Controls & Prevents: For treatment and control of susceptible tapeworms, roundworms (including arterial larval stages of *Strongylus vulgaris* and benzimidazole resistant small strongyles), bots and skin lesions caused by *Habronema* and *Draschiaspp.* (summer sores) and *Onchocercaspp.* microfilariae (cutaneous onchocerciasis).

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure


[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)



Custom Product Download
Missing Attachment
Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

140.3mg/g Praziquantel, 18.7g/kg Ivermectin. For treatment and control of susceptible tapeworms, roundworms, bots and skin lesions caused by summer sores. 1 x 700kg Horse, 7.49g.

Controls & Prevents: Treats and controls tapeworm, large strongyles, small strongyles (including strains that have become resistant to benzimidazole), pinworms, ascarids (roundworms), hairworms, large-mouth stomach worms, neck threadworms, bots, lungworms, intestinal threadworms, Habronema and Draschia spp. cutaneous larvae (summer sores) and Onchocerca spp. microfilariae (cutaneous onchocerciasis).

Dose Rates: The contents of one syringe will treat one horse of 700kg bodyweight. Each weight marking on the syringe plunger will deliver 1.07g of paste which is sufficient to treat 100kg bodyweight.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure



[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Low volume pastes presented in ergonomic syringes for easy administration. Less chance of spitting out, and syringe ideal for smaller hands.

18.7g/kg Ivermectin. 1 x 700kg Horse. 7.49g.

Controls & Prevents: Effective against Benzimidazole (BZ) resistant worms (small strongyles) – some strains of bloodworms have become resistant to BZ products, EQUIGARD can control BZ resistant worms

Dose Rates: The contents of one syringe will treat one horse of 700kg bodyweight. Each weight marking on the syringe plunger will deliver 1.07g of paste which is sufficient to treat 100kg bodyweight.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure



[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Containing Eprinomectin, one of the world's most potent endectocides, iO Eprinomax provides broad-spectrum control of internal and external parasites with the flexibility of Nil meat and Nil milk withholding periods.

Active Constituent: 5 mg/ml Eprinomectin

Controls & Prevents: For the treatment and control of gastrointestinal roundworms (including inhibited immature *Ostertagia ostertagi*) and lungworm in cattle. For sucking and biting lice, sarcoptic and chorioptic mange mites, and buffalo fly in beef and dairy cattle. iO EPRINOMAX Pour-On aids in control of cattle tick (*Rhipicephalus microplus*). For the treatment and control of gastrointestinal roundworms and lungworm in deer.

Recommended Applicators: Large cattle (300kg +): iO Pour-on applicator 65ml. Smaller cattle (up to 300kg): iO Drench/Pour-On applicator 30mL.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)



Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Rapidly kills and destroys lice very soon after treatment.

Active Constituent: 350g/L Temephos. Solvent: Liquid hydrocarbons 492 g/L

Made in Australia.

Dose Rates & Administration Instructions: Yard sheep the night before dipping. Ensure adequate water is available. Preferably dip early in the morning and allow sheep to dry before nightfall. No need to reinforce the dip as the dip is non stripping making it really a simple application. Can be applied to wet or dry sheep.

Store below 30 degrees.

Dipping Management The sheep should be thoroughly wetted. Sheep lice live on the skin of the sheep and complete saturation of the fleece is necessary. In a plunge dip each sheep should swim for at least 30 seconds and have its head plunged under twice. Shower dips should be checked prior to commencement of dipping to ensure that the pump and nozzles are operating at the correct pressure and volumes as stipulated by the manufacturer.



Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent 10mg/ml Abamectin, 200mg/ml Levamisole

Dose Rates SURESHOT DUO POUR-ON dose rate is 1 mL/20 kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline of the animal, using the SURESHOT DUO POUR-ON applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected. Only use the pour-on applicator supplied for use with SURESHOT DUO POUR-ON. SURESHOT DUO POUR-ON is not compatible with other guns. After use, rinse gun with warm soapy water.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure


[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)



Custom Product Download
Missing Attachment
Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Levistar LV contains levamisole, a member of the imidazothiazole (clear) group of anthelmintics (drenches) and is effective in cattle and sheep against: mature and immature Barber's Pole Worm, Stomach Hair Worm, Black Scour Worm, Small Intestinal worm (*Cooperia* spp.), Thin Necked Intestinal Worm, Hookworm,

Large Mouthed Bowel Worm, *Strongyloides*, Nodule and Large Bowel Worms, Large Lungworm and Small Brown Stomach Worm in sheep and mature (adult only) *Ostertagia* spp. in cattle under 12 months of age. Where *Ostertagia* are the problem in cattle over 12 months of age, Starmec Pour-On for Beef and Dairy Cattle is the recommended product.

Controls both susceptible and benzimidazole resistant strains of Barber's pole worm (*Haemonchus* species) and Black scour worm (*Trichostrongylus* species) in sheep.

Active constituents: 80 g/L Levamisole Hydrochloride (equivalent to 68 g/L Levamisole)

Dosing and Administration: Levistar LV is given as a drench by mouth in cattle and sheep.

Withholding periods:



Meat: DO NOT USE less than 3 days before slaughter for human consumption.

Milk: Zero (0) days.

Trade Advice: Export Slaughter Interval (ESI): DO NOT USE less than 7 days before slaughter for export.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Pack size: Contains 5 x 500 g measure packs which it is illegal to sell separately.

Administration Instructions: This pack dissolves on contact with water. Open outer sachet. To dissolve in water, simply tip inner soluble pack directly into bucket and stir. Dispense as per mixing instructions advised on label.

Controls & Prevents:

ROUTINE TICK CONTROL (NSW, QLD, VIC, NT only): For routine control of specific types of ticks on cattle, apply HRC EXITRAZ at the following intervals:

CATTLE TICK (*Rhipicephalus* [*Boophilus*] *microplus*): intervals of 19 to 21 days.

BUSH TICK (NZ CATTLE TICK) (*Haemaphysalis longicornis*): intervals of 7 to 21 days.

PARALYSIS TICK (*Ixodes holocyclus*): intervals of 7 to 10 days. DO NOT leave longer intervals between treatments when ticks are active. For further information consult your Stock Inspector, District Veterinary Officer or Tick Extension Officer.

Breed Category

Cattle



Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Can be safely administered to: All classes of sheep across Australia

Active Constituents: 82.5 g/L LEVAMISOLE HYDROCHLORIDE and 36 g/L ALBENDAZOLE OXIDE

Withholding Periods:

- MEAT: Off Shears: DO NOT USE less than 10 days before slaughter for human consumption.
- MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.
- Export Slaughter Interval (ESI): Do not used less than 10 day before slaughter for export.
- Please refer to label and SDS for further information.

Safety Directions:

Poisonous if swallowed. Harmful if absorbed by skin. May irritate eyes and skin. Avoid contact with eyes and skin. Wear rubber gloves. Wash hands after use.

Storage Instructions:



Store below 30oC (room temperature) in a dry place in tightly closed in original container.

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 5 mg/mL Doramectin

Application: 30ml Variable Applicator with Pour-On Nozzle or Pour-On Applicator 65ml.

Dosing & Administration: Apply topically along the mid-line of the back in a narrow strip between the withers and tailhead.

Use the dose rate, according to the weight of the animal, given in the tables below. The dose rate should be based on the heaviest member in the weight range (given under the heading "weight (kg)"). Cattle should be weighed with either scales or a weighband.

Bodyweight (kg) Dose (mL)

40–50	5
51–75	7.5
76–100	10
101–150	15
151–200	20

Bodyweight (kg) Dose (mL)

201–250	25
251–300	30
301–350	35
351–400	40
401–450	45
451–500	50
501–550	55
551–600	60
601–650	65

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 300 MG/ML Triclabendazole & 5 MG/ML Abamectin

Available in the following sizes: 1L, 2.5L, 5L,15L

Dosing & Administration:

HRC Abatech Ultra Pour On dose rate is 1 mL/10 kg body weight.

Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.

Liveweight (kg)	Dose Vol (mL)	No. of Cattle Treated	
1L	2.5 L	5 L	15 L
51 – 75	7.5	133	333 666 2000
76 – 100	10	100	250 500 1500
101 – 125	12.5	80	200 400 1200

126 – 150	15	66	166	333	1000
151 – 175	17.5	57	142	285	857
176 – 200	20	50	125	250	750
201 – 250	25	40	100	200	600
251 – 300	30	33	83	166	500
301 – 350	35	28	71	142	428
351 – 400	40	25	62	125	375
401 – 450	45	22	55	111	333
451 – 500	50	20	50	100	300
501 – 550	55	18	45	90	272
551 – 600	60	16	41	83	250
601 – 650	65	15	38	76	230

For animals heavier than 650 kg, dose at a rate of 1 mL/10 kg body weight.

A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Abatech Ultra Pour On is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)



Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Abastar Plus Selenium contains abamectin, a member of the macrocyclic 'ML' family of drenches and also contains selenium which is an aid in control and treatment of selenium deficiency in sheep and lambs. Abastar Plus Selenium is effective against abamectin sensitive strains of the following adult and immature internal parasites of sheep. **Active Constituent:** 0.8 g/L Abamectin. 0.4 g/L Selenium (as Sodium Selenate) **Controls & Prevents Adult and immature gastrointestinal roundworms such as:**

- *Haemonchus contortus* – Barber's pole worm (including inhibited L4 stage)
- *Haemonchus placei* – Large stomach worm
- *Ostertagia circumcincta* – Small brown stomach worm (including inhibited L4 stage)
- *Trichostrongylus axei* – Stomach hair worm
- *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus* – Black scour worm
- *Cooperia curticei*, *Cooperia oncophora* – Small intestinal worm
- *Nematodirus spathiger*, *Nematodirus filicollis* – Thin necked intestinal worm
- *Chabertia ovina* – Large mouthed bowel worm
- *Oesophagostomum columbianum* – Nodule worm
- *Oesophagostomum venulosum* – Large bowel worm
- *Trichuris ovis* – Whipworm

- *Strongyloides papillosus* – Intestinal threadworm

Lungworm:

- *Dictyocaulus filaria* – Large lungworm
- *Oestrus ovis* – Nasal bot
- *Psorergates ovis* – Itch mite

Dosage and Administration: Abastar Plus Selenium dose rate is 1 mL/4 kg bodyweight.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active constituents: 2.0 G/L Abamectin, 1.0 G/L Selenium (as sodium selenate)

Target parasites: Abastar Plus Selenium LV contains abamectin, a member of the avermectin family of drenches. It is effective against abamectin sensitive strains of the following internal parasites:

Adult and immature gastrointestinal roundworms:

Barber's pole worm – *Haemonchus contortus*

Small brown stomach worms – *Ostertagia* spp.

Black scour worms – *Trichostrongylus* spp.

Small intestinal worms – *Cooperia* spp.

Thin-necked intestinal worms – *Nematodirus* spp.

Large bowel worm – *Oesophagostomum venulosum*

Lungworm: *Dictyocaulus filaria*

For the control of Sheep Nasal Bot – *Oestrus ovis*

Recommended Applicator: 30mL Variable Applicator

Withholding periods:

Meat: DO NOT USE less than 14 days before slaughter for human consumption.

Milk: DO NOT USE in lactating ewes where milk may be used or processed for human consumption

Trade Advice – Export Slaughter Interval (ESI): An ESI has not been established for this product.

Dosage & administration: 1mL HRC Abastar Plus Selenium LV per 10kg bodyweight.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Features:

- Wide safety margin when administered to cattle & pigs.
- Low irritant formulation, which will minimise any potential reaction to the injection site post administration.
- Broad spectrum treatment for the control of gastrointestinal roundworms, lungworms, sucking lice, biting lice, cattle tick and mange mites.
- The most reliable and accurate dosage format, which ensures animals receive the optimal dose.

Active Constituent: 10 mg/mL Doramectin

Application: Metal Drencher 20mL with Injector Conversion Kit.

Dosing & Administration: Use contents within 28 days of first broaching of the vial. Store upright. Discard the unused portion. Use a multi-dose syringe for multiple treatments to minimise the number of broachings.

CATTLE: 1 mL per 50 kg of bodyweight SC

PIGS: 1 mL per 33 kg (0.3 mL per 10 kg) of bodyweight IM



Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Nil meat and milk withholding provides flexibility particularly if clearing cattle of ticks prior to sale. Following treatment, cattle will be clean of ticks within 3-4 days, with larval ticks prevented from attaching for up to 7 days post treatment.

Active Constituent: 125 g/L Amitraz

Available in the following sizes: 400ML & 2L

Application: Coarse Spray with pump

Dosing & Administration

LIVESTOCK	PEST	STATE	RATE	CRITICAL COMMENTS
Cattle	Cattle Tick [Rhipicephalus (Boophilus) microplus]	NSW, QLD, WA and NT only	400mL per 200L water	Treat at intervals of 19-21 days Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick.Refer also to General Directions
	Paralysis Tick (Ixodes holocyclus)	NSW, QLD, WA and NT only	400mL per 200L water	Treat at intervals of 7-10 days Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick.Refer also to General Directions
	Bush Tick (NZ Cattle Tick) (Haemaphysalis longicornis)	NSW, QLD, WA, NT and VIC	400mL per 200L water	Treat at intervals of 7-21 days Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick.Refer also to General Directions
Sheep, goats, deer and certain circus animals	Cattle Tick [Rhipicephalus(Boophilus) microplus]	NSW only	400mL per 200L water	Treat animals on quarantined holdings at 14 to 21 day intervals. Treat animals being moved at 3 to 7 day intervals.
Pigs	Mange (Sarcoptes scabiei var. suis)	All States	40 mL per 10L water	Remove feed and bedding from pens, cover drinking bowls and clean out pen. Spray pigs with minimum 2L spray wash, especially inside ears and legs, under jowls and areas covered by scabs. Replace discarded bedding with clean material. Repeat treatment after 7-10 days. Refer also to General Directions.

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure



[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Designed for use where resistance to single actives may be present, Tripletec HV supports effective parasite control in modern sheep systems. By combining actives from different chemical groups, it helps address worm populations that may not be adequately controlled by single-active drenches alone.

Tripletec HV is effective against susceptible, single and dual resistant parasite strains and also provides control of additional parasites including lungworm, nasal bot (larval stages) and itch mite.

Tripletec HV also includes selenium and cobalt to assist with trace element supplementation in sheep grazing in areas where these minerals may be deficient.




Active Ingredients

Levamisole (as hydrochloride) 25.5 g/L, Albendazole 20 g/L, Cobalt 1.76 g/L (as EDTA), Abamectin 0.8 g/L, Selenium 0.4 g/L (as sodium selenate)

Key Benefits

- Triple-active, 3-way combination drench
- Broad internal parasite control for sheep
- Suitable for use where resistance to single actives may be present
- Supports integrated parasite management programs
- Effective against susceptible, single and dual resistant strains

- 
- Controls additional parasites including lungworm, nasal bot (larval stages) and itch mite
 - Includes selenium and cobalt for use in deficient areas

Effective Against

- Sheep – Internal parasites, including:
 - Barber's Pole Worm
 - Black Scour Worm
 - Small Brown Stomach Worm
 - Cooperia spp.

- 
- Nematodirus spp.

 - Lungworm, nasal bot (larvae) and itch mite

Dosage

Refer to the label for further information

Withholding Periods

Meat: DO NOT USE less than 14 days before slaughter for human consumption.

Export Slaughter Interval (ESI)



DO NOT USE less than 42 days before slaughter for export.

Before using this product, confirm the current ESI from the distributor on (03) 5820 8400 or the APVMA website (apvma.gov.au/residues).

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Dose Rates

The recommended dose rate for iO Mectec LV is 1 mL/20kg body weight. Apply along the topline of the animal.

Applicator


iO 30ml Variable Applicator – Pour-On Nozzle

Withholding periods

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

MILK: Zero (0) days.

ESI: DO NOT use less than 42 days before slaughter for export.



Storage

Store below 30°C (room temperature).

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Controls & Prevents: For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*) and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle. For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses. Provides protection from the Biting Midge (*Culicoides brevitarsus*) for at least 14 days with effective control of blood feeding *C. brevitarsus* for 6 days post treatment.

Active Constituent: 15.0g/L Deltamethrin

Dose Rates for Cattle: 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice (<i>Bovicola bovis</i>) (<i>Lignognathus vituli</i>) (<i>Haematopinus eurysternus</i>)	5 mL 15mL 301-400kg	Up to 100kg 10mL 201-300kg 20mL 401-500kg	101-200kg Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>) House Fly (<i>Musca domestica</i>) Biting Midge (<i>Culicoides sp.</i>)	30mL apply 5mL per 100kg liveweight	501-600kg For cattle over 600kg	Apply to the whole herd – retreatment may be necessary.

Dose Rate for Horses: 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>)	5 mL 101-200kg 15mL 201-300kg 20mL 301-400kg 25mL 401-500kg 30mL 501-600kg For horses over 600kg apply 5mL per 100kg liveweight	Up to 100kg 10mL 201-300kg 401-500kg 501-600kg For horses over 600kg	Using the correct measured dose, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump. When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.



Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Quick Facts & Benefits: Protects sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.

May be used in conjunction with worm control drenches and at the same time as vaccination.

Contains a blue dye to identify treated sheep. This dye is scourable from the wool. Is safe for use on breeding rams.

Proudly Australian Made & Owned.

Active Constituent: 35 g/L Imidacloprid Also contains: 300 g/L N-Methyl-2-Pyrrolidone (solvent)

Dose Rates & Administration Instructions: Refer to the dosage table on label. One 20L pack will treat 333 sheep & one 5L pack will treat 83 sheep (based on 30.1 – 55kg body weight).



Recommended Applicators: Apply product only with the 6-stripe applicator.

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 100 mg/g Praziquantel & 8 mg/g Ivermectin.

Double strength formulation of ivermectin and praziquantel.

Dose Rates: 30ml syringe treats 2 x 600kg horses. Easy to administer. Wide safety margin.

Controls & Prevents: For the treatment and control of roundworms (including arterial larval stages of *Strongylus vulgaris* and benzimidazole resistant small strongyles), tapeworms, bots and skin lesions (summer sores) caused by *Habronema* and *Draschia* spp. and *Onchocerca* spp. microfilariae (cutaneous onchocerciasis).

Breed Category



Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

With persistent activity against internal parasites for up to 120 days (see below table for details) or cattle/tick for at least 51 days, no other product offers longer term protection ensuring optimum health, performance and fertility.

OXISTAR LONG ACTING INJECTION FOR CATTLE is effective against sensitive strains of the following parasites:

Internal parasites: Mature (adult) and immature (L4): Barber's pole worm, *Ostertagia ostertagi* / *Ostertagia lyrata* (including inhibited larvae), Black scour worms, Small intestinal worm, Nodule worm, Hookworm, Whipworm, Lungworm and Adult nematodes.

External parasites: Lice (Sucking lice). Aids in the control of *Bovicola bovis* (biting lice), Mites and Ticks.



Dose Rates 1 mg moxidectin/kg liveweight (1 mL/100 kg liveweight)

Breed Category
Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

Active Constituent: 200 g/L Chlorfenvinphos (an anticholinesterase compound)

Application: Backrubber or overspray.

Dosing & Administration:

Pest: Buffalo fly (*Haematobia irirritans exigua*)

Treatment: Application by backrubber. Make backrubber available to cattle each day and all day during the buffalo fly season.



State: QLD, WA, NSW, NT only

Rate: Mix 1 L EXIGUARD Buffalo Fly Insecticide per 20 L of new, non-recycled motor oil or clean mineral oil. Charge backrubber with 20 mL treated new, non-recycled motor oil or clean mineral oil per animal every three weeks. Do not exceed this rate.

Critical comments: Refer to mixing instructions on the label.

NOT TO BE USED ON ANIMALS PRODUCING MILK FOR HUMAN CONSUMPTION WHEN TREATMENT IS BY BACKRUBBER APPLICATION.”

Breed Category

Cattle

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1