



Starvec Injection will treat and control internal parasites including gastrointestinal roundworms, lungworms, eyeworms, sucking and biting lice, mites and cattle tick in cattle.

**Active Constituent:** 10mg/ml Ivermectin

**Available in the following sizes:** 500ML

**Applicator:** 20ml Injector (Drencher with conversion kit)

**Dosing & Administration:** Use the contents within 24 hours of first broaching of the vial. Discard the unused portion.

The recommended dose is 1 mL STARMEC Ivermectin Injection for Cattle per 50kg liveweight (equivalent to 200 µg ivermectin per kg liveweight) by subcutaneous injection. DO NOT administer by the intravenous or intramuscular route.

**Liveweight (kg) Dose volume (mL) Treatments per 500mL**

Up to 100	2	250
101 – 150	3	166
151 – 200	4	125
201 – 250	5	100
251 – 300	6	83
301 – 350	7	71
351 – 400	8	62
401 – 450	9	55
451 – 500	10	50
501 – 550	11	45
551 – 600	12	41
601 – 650	13	38



Breed Category

Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Active constituent:** 20 g/L SPINOSAD.

**Recommended applicator:** iO 30ml Applicator with 4 hole T Bar Kit (31845).

**Dosage & Administration:** Apply iO SPINOFF POUR-ON evenly along the backline from the poll to the base of the tail using a manual or powered applicator with an iO SPINOFF POUR-ON 4-hole t-bar nozzle. Halve the dose volume and apply in 2 bands, approximately 5 cm from the mid-point of the backline on each side. Applying the product to the flanks will improve the spread of iO SPINOFF POUR-ON.

### Breed Category

Cattle

### Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

### Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

### Safety Data Sheet



[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

---

**Active Constituent:** 5 g/L Moxidectin

**Application:** 30ml Variable Applicator with Pour-On Nozzle or Pour-On Applicator 65ml.

**Dosing & Administration:** MOXISTAR CATTLE POUR-ON should be applied along the top of the back from withers to the base of the tail. The volume to use is 1 mL per 10 kg liveweight. This will administer the correct dose of 0.5 mg/kg liveweight.

MOXISTAR CATTLE POUR-ON can be used in lactating dairy cattle, but DO NOT RETREAT LACTATING CATTLE AT LESS THAN 21 DAY INTERVALS.

**Liveweight (kg) Dose (mL)**

31-40	4
41-50	5
51-60	6
61-70	7
71-80	8

Liveweight (kg) Dose (mL)

81-90	9
91-100	10
101-125	12.5
126-150	15
151-175	17.5
176-200	20
201-250	25
251-300	30
301-350	35
351-400	40
401-450	45
451-500	50
501-550	55

Breed Category

Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label




[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

For the treatment and control of moxidectin sensitive gastrointestinal parasites (including benzimidazole and/or levamisole resistant strains), lungworm and itch mite of sheep. An aid in the control of selenium responsive conditions.

Moxistar LV is highly effective against gastrointestinal parasites including Barber's Pole worm, Black scour worm, Stomach hair worm, Small brown stomach worm, Small intestinal worm, Thin necked intestinal worm, Intestinal threadworm, Nodule worm, Large mouthed bowel worm, Large lungworm & itch mite.

**Active Constituent:** 2 g/L Moxidectin 1 g/L Selenium (as Sodium selenate)

**Dosage and Administration:** MOXISTAR LV Se is ready to use. Administer 1 mL per 10 kg liveweight using any standard drenching equipment to give a dose of 0.2 mg moxidectin/kg and 0.1mg selenium/kg liveweight. Check dose rates and equipment before drenching commences.

### Withholding Periods:

- MEAT: DO NOT USE less than 7 days before slaughter for human consumption.
- MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

- EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export.

Breed Category

Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

Containing moxidectin in a long-acting formulation, it provides sustained control of internal parasites in sheep during periods of challenge.

**Active Ingredient:** 20mg/ml Moxidectin

### Benefits

- Provides control of internal parasites in sheep
- Contains moxidectin, a highly potent member of the macrocyclic lactone (ML) family
- Effective against parasites susceptible to ML drenches, including gastrointestinal roundworms and large lungworm
- ≥91 days persistent activity against Barber's Pole worm and Small Brown Stomach worm
- Up to 49 days persistent activity against Black Scour worm
- Up to 4 months protection during severe Barber's Pole worm challenge
- Long duration of activity may reduce the need for frequent drenching
- Suitable for use in highly contaminated pastures
- Controls nasal bot (larval stages) and itch mite
- Simple subcutaneous injection for ease of application

- Wide margin of safety when used as directed

When administered at the recommended dose rate, Moxistar LA Sheep Injection provides:

- At least 91 days persistent activity against Barber's Pole worm (*Haemonchus contortus*)
- At least 91 days persistent activity against Small Brown Stomach worm (*Teladorsagia circumcincta*)
- Up to 49 days persistent activity against Black Scour worm (*Trichostrongylus* spp.)
- During periods of severe Barber's Pole worm challenge, protection may extend for up to 4 months when used as directed.
- Provides long-acting control of key internal parasites in sheep, including extended protection against barber's pole worm and small brown stomach worm.
- Delivers persistent activity against black scour worm and helps protect sheep under heavy parasite challenge.

#### Withholding Period/ESI

- Meat: 91 days
- Milk: Do not use in female sheep which are producing, or may in the future produce, milk or milk products for human consumption
- ESI: 133 days

Breed Category

Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label




[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

iO Tick Off Duo contains fluazuron and ivermectin to give the dual benefits of tick control (fluazuron) and broad spectrum internal/external parasite protection (ivermectin) in one easy to use pour-on. Use in strategic cattle tick programs when broad spectrum parasite control is required in tick infested geographies.

### Active Constituent:

15 g/L Fluazuron, 5 g/L Ivermectin

**Controls & Prevents:** Cattle Tick (*Rhipicephalus microplus*), ivermectin-sensitive gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, sucking and biting lice, sarcoptic and chorioptic mites, and buffalo fly (*Haematobia irritans exigua*) on Beef Cattle.

### Breed Category

Cattle

### Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

### Buy Now Tuckers



<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Active Constituent:** 5 g/L Ivermectin.

**Controls & Prevents:** For the treatment and control of ivermectin sensitive:

Gastrointestinal nematodes:

Adult and immature (L4): Small Brown Stomach Worm (including inhibited stages) (*Ostertagia ostertagi*), Barber's Pole Worm (*Haemonchus placei*), Stomach Hair Worm (*Trichostrongylus axei*), Intestinal Hair Worm (*T. colubriformis*), Small Intestinal Worm (*Cooperia* spp.), Nodule Worm (*Oesophagostomum radiatum*); and Adult stages of *Cooperia oncophora*, *C. punctata*, *Oesophagostomum venulosum*, Whip Worm (*Trichuris* spp.), Intestinal Threadworm (*Strongyloides papillosis*); and Immature stages (L4) of Thin Necked Intestinal Worm (*Nematodirus* spp.).


Lungworms: (adult and immature) *Dictyocaulus viviparus*.

Eyeworm: *Thelazia* spp. (adult stages).

Sucking and Biting Lice: *Linognathus vituli*, *Haematopinus euryesternus*, *Bovicola bovis*, *Solenopotes capillatus*.

Mites: *Chorioptes bovis*, *Sarcoptes scabiei* var *bovis*.

Buffalo Flies: *Haematobia irritans exigua* g. Cattle Ticks: (*Boophilus microplus*) including organophosphate, synthetic pyrethroid- and amidine-resistant strains.



**Administration Instructions:** This product is formulated only for external application to cattle; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead.

**Dose Rates** The dose is 1mL for each 10kg of liveweight. Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers etc.).

When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid overdosing. Do not underdose. Cattle should be weighed with either scales or with a weighband.

Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Active constituents:** Ivermectin 10g/L & Clorsulon 100g/L

Mectec Plus is a ready to use broad-spectrum endectocide injection for cattle containing Ivermectin and Clorsulon for the treatment and control of Ivermectin and Clorsulon sensitive strains of internal and external parasites of cattle, including adult liver flukes.

The recommended dose rate is 1mL/50kg live weight by subcutaneous injection. Heavier animals (for example mature bulls) should receive an additional 1 mL for each 50kg over 650kg.

Liveweight (kg)	Dose (mL)	No. of animals treated
Up to 100	2	250
101 – 150	3	166
151 – 200	4	125
201 – 250	5	100
251 – 300	6	83

Liveweight (kg)	Dose (mL)	No. of animals treated
301 – 350	7	71
351 – 400	8	62
401 – 450	9	55
451 – 500	10	50
501- 550	11	45
551 – 600	12	41
601 – 650	13	38

**Withholding periods:**

**MEAT:** DO NOT USE less than 28 days before slaughter for human consumption.

**MILK:** Zero (0) days

**EXPORT SLAUGHTER INTERVAL (ESI):** DO NOT slaughter for export less than 42 days after treatment.

**Recommended Applicators:** iO 15 mL Injector

**Storage:** Store below 30c (Room temperature) and protect from light in a closed, original container.

Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers



<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Active Constituent:** 16g/L Ivermectin

**Dose Rates & Administration Instructions:** JETTING FOR BLOWFLY STRIKE: When applied as directed by hand jetting or by modified Harrington jetting race JET AWAY BLOWFLY AND LICE JETTING FLUID FOR SHEEP protects sheep against strike by blowflies (*Lucilia cuprina*, *L. sericata*, *Chrysomya rufifacies* and *Calliphora nociva*) (including OP resistant strains of *Lucilia cuprina*) for 12 weeks under low to moderate fly pressure and will control maggots in existing strikes in the treated areas.

HAND DRESSING OF STRUCK SHEEP: JET AWAY can be applied at the blowfly dilution rate for treatment of blowfly strike wounds. Sheep should be sheared/clipped around the strike wound before saturating the area.

HAND JETTING FOR LICE: When applied as directed by hand-jetting JET AWAY also treats biting lice (*Bovicola ovis*), including synthetic pyrethroid resistant strains in long woolled sheep until the next shearing. Testing has confirmed that JET AWAY mixes with waters of various hardness. Use either a jetting handpiece with multiple nozzles that can be combed through the fleece or a "Dutjet" jetting wand. The jetting handpiece should deliver a penetrating jet of fluid without producing a spray or mist. A modified Harrington jetting race can also be used.



Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

Containing Eprinomectin, one of the world's most potent endectocides, iO Eprinomax provides broad-spectrum control of internal and external parasites with the flexibility of Nil meat and Nil milk withholding periods.

**Active Constituent:** 5 mg/ml Eprinomectin

**Controls & Prevents:** For the treatment and control of gastrointestinal roundworms (including inhibited immature *Ostertagia ostertagi*) and lungworm in cattle. For sucking and biting lice, sarcoptic and chorioptic mange mites, and buffalo fly in beef and dairy cattle. iO EPRINOMAX Pour-On aids in control of cattle tick (*Rhipicephalus microplus*). For the treatment and control of gastrointestinal roundworms and lungworm in deer.

**Recommended Applicators:** Large cattle (300kg +): iO Pour-on applicator 65ml. Smaller cattle (up to 300kg): iO Drench/Pour-On applicator 30mL.

### Breed Category

Cattle

### Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>



Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

Rapidly kills and destroys lice very soon after treatment.

**Active Constituent:** 350g/L Temephos. Solvent: Liquid hydrocarbons 492 g/L

Made in Australia.

**Dose Rates & Administration Instructions:** Yard sheep the night before dipping. Ensure adequate water is available. Preferably dip early in the morning and allow sheep to dry before nightfall. No need to reinforce the dip as the dip is non stripping making it really a simple application. Can be applied to wet or dry sheep.

Store below 30 degrees.

**Dipping Management** The sheep should be thoroughly wetted. Sheep lice live on the skin of the sheep and complete saturation of the fleece is necessary. In a plunge dip each sheep should swim for at least 30 seconds and have its head plunged under twice. Shower dips should be checked prior to commencement of dipping to ensure that the pump and nozzles are operating at the correct pressure and volumes as stipulated by the manufacturer.



Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery



### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Active Constituent** 10mg/ml Abamectin, 200mg/ml Levamisole

**Dose Rates** SURESHOT DUO POUR-ON dose rate is 1 mL/20 kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline of the animal, using the SURESHOT DUO POUR-ON applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected. Only use the pour-on applicator supplied for use with SURESHOT DUO POUR-ON. SURESHOT DUO POUR-ON is not compatible with other guns. After use, rinse gun with warm soapy water.

### Breed Category

Cattle

### Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

### Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

### Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)



Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

Active Constituent

10mg/ml Ivermectin

Product Sizes

500ml

Made in Australia

1

---

**Active Constituent:** 5 mg/mL Doramectin

**Application:** 30ml Variable Applicator with Pour-On Nozzle or Pour-On Applicator 65ml.

**Dosing & Administration:** Apply topically along the mid-line of the back in a narrow strip between the withers and tailhead.

Use the dose rate, according to the weight of the animal, given in the tables below. The dose rate should be based on the heaviest member in the weight range (given under the heading "weight (kg)"). Cattle should be weighed with either scales or a weighband.

**Bodyweight (kg) Dose (mL)**

40–50	5
51–75	7.5
76–100	10
101–150	15
151–200	20

Bodyweight (kg) Dose (mL)

201–250	25
251–300	30
301–350	35
351–400	40
401–450	45
451–500	50
501–550	55
551–600	60
601–650	65

Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

### Features:

- Wide safety margin when administered to cattle & pigs.
- Low irritant formulation, which will minimise any potential reaction to the injection site post administration.
- Broad spectrum treatment for the control of gastrointestinal roundworms, lungworms, sucking lice, biting lice, cattle tick and mange mites.
- The most reliable and accurate dosage format, which ensures animals receive the optimal dose.

**Active Constituent:** 10 mg/mL Doramectin

**Application:** Metal Drencher 20mL with Injector Conversion Kit.

**Dosing & Administration:** Use contents within 28 days of first broaching of the vial. Store upright. Discard the unused portion. Use a multi-dose syringe for multiple treatments to minimise the number of broachings.

**CATTLE:** 1 mL per 50 kg of bodyweight SC

**PIGS:** 1 mL per 33 kg (0.3 mL per 10 kg) of bodyweight IM



Breed Category

Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Active Constituent:** 300 MG/ML Triclabendazole & 5 MG/ML Abamectin

Available in the following sizes: 1L, 2.5L, 5L,15L

### Dosing & Administration:

HRC Abatech Ultra Pour On dose rate is 1 mL/10 kg body weight.

Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.

### Liveweight (kg) Dose Vol (mL) No. of Cattle Treated

1L	2.5 L	5 L	15 L
51 – 75	7.5	133	333 666 2000
76 – 100	10	100	250 500 1500
101 – 125	12.5	80	200 400 1200

126 – 150	15	66	166	333	1000
151 – 175	17.5	57	142	285	857
176 – 200	20	50	125	250	750
201 – 250	25	40	100	200	600
251 – 300	30	33	83	166	500
301 – 350	35	28	71	142	428
351 – 400	40	25	62	125	375
401 – 450	45	22	55	111	333
451 – 500	50	20	50	100	300
501 – 550	55	18	45	90	272
551 – 600	60	16	41	83	250
601 – 650	65	15	38	76	230

For animals heavier than 650 kg, dose at a rate of 1 mL/10 kg body weight.

A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Abatech Ultra Pour On is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product.

Breed Category

Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>



Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

### Dose Rates

The recommended dose rate for iO Mectec LV is 1 mL/20kg body weight. Apply along the topline of the animal.

### Applicator

iO 30ml Variable Applicator – Pour-On Nozzle

### Withholding periods

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

MILK: Zero (0) days.

ESI: DO NOT use less than 42 days before slaughter for export.

## Storage

Store below 30°C (room temperature).

Breed Category

Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Controls & Prevents:** For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*) and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses. Provides protection from the Biting Midge (*Culicoides brevitarsus*) for at least 14 days with effective control of blood feeding *C. brevitarsus* for 6 days post treatment.

**Active Constituent:** 15.0g/L Deltamethrin

**Dose Rates for Cattle:** 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice ( <i>Bovicola bovis</i> ) ( <i>Lignognathus vituli</i> ) ( <i>Haematopinus eurysternus</i> )	5 mL 15mL 301-400kg	Up to 100kg 10mL 201-300kg 20mL 401-500kg	Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
Buffalo Fly ( <i>Haematobia irritans exigua</i> ) Stable Fly ( <i>Stomoxys calcitrans</i> ) House Fly ( <i>Musca domestica</i> ) Biting Midge ( <i>Culicoides sp.</i> )	30mL apply 5mL per 100kg liveweight	501-600kg For cattle over 600kg	Apply to the whole herd – retreatment may be necessary.

Dose Rate for Horses: 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Buffalo Fly ( <i>Haematobia irritans exigua</i> ) Stable Fly ( <i>Stomoxys calcitrans</i> )	5 mL 101-200kg 15mL 20mL 401-500kg 30mL horses over 600kg apply 5mL per 100kg liveweight	Up to 100kg 10mL 201-300kg 301-400kg 25mL 501-600kg For	Using the correct measured dose, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump. When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.



Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)

Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

**Quick Facts & Benefits:** Protects sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.

May be used in conjunction with worm control drenches and at the same time as vaccination.

Contains a blue dye to identify treated sheep. This dye is scourable from the wool. Is safe for use on breeding rams.

Proudly Australian Made & Owned.

**Active Constituent:** 35 g/L Imidacloprid Also contains: 300 g/L N-Methyl-2-Pyrrolidone (solvent)

**Dose Rates & Administration Instructions:** Refer to the dosage table on label. One 20L pack will treat 333 sheep & one 5L pack will treat 83 sheep (based on 30.1 – 55kg body weight).



**Recommended Applicators:** Apply product only with the 6-stripe applicator.

Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1

---

With persistent activity against internal parasites for up to 120 days (see below table for details) or cattle/tick for at least 51 days, no other product offers longer term protection ensuring optimum health, performance and fertility.

OXISTAR LONG ACTING INJECTION FOR CATTLE is effective against sensitive strains of the following parasites:

Internal parasites: Mature (adult) and immature (L4): Barber's pole worm, *Ostertagia ostertagi* / *Ostertagia lyrata* (including inhibited larvae), Black scour worms, Small intestinal worm, Nodule worm, Hookworm, Whipworm, Lungworm and Adult nematodes.

External parasites: Lice (Sucking lice). Aids in the control of *Bovicola bovis* (biting lice), Mites and Ticks.



**Dose Rates** 1 mg moxidectin/kg liveweight (1 mL/100 kg liveweight)

Breed Category  
Cattle

Buy Now AIRR

<https://airr.com.au/product/livestock/cattle/cattle-drenches/starmec-cattle-injectable-500ml-ivermectin/>

Buy Now Tuckers

<https://tuckersstore.com.au/product/starmec-cattle-injectable-500ml-ivermectin/>

Safety Data Sheet

[Starmec-Ivermectin-Injection-for-Cattle-84437-SDS-valid-Aug-27](#)

Product Brochure

[Starmec-Injectable-A4-Flyer---SCREEN](#)


Product Label

[Starmec-Inj-for-Cattle---Label-80x130mm---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery





### Short Description

Convenient one shot Ivermectin injection for control of ivermectin sensitive internal and external parasites in cattle. For those producers which prefer injections vs pour-ons, Starmec injection will do the job.

### Active Constituent

10mg/ml Ivermectin

### Product Sizes

500ml

Made in Australia

1