



iO Tick Off is a ready-to-use pour-on formulation. iO Tick Off breaks the tick life cycle. When treated with iO Tick Off, immature ticks are unable to molt to the next stage of their life cycle, and no larvae will hatch from eggs laid by treated females.

Active Constituent:

25g/L Fluazuron

Solvents:

100 g/L 1-dodecyl-2-pyrrolidone

100 g/L 1-octyl-2-pyrrolidone

415 g/L 1-methyl-2-pyrrolidone

Controls & Prevents: For strategic control of Cattle Tick, *Boophilu microplus*, (including all known acaricide resistant strains) on Beef Cattle.

Dose Rates & Administration Instructions: Apply iO Tick Off in two bands, roughly 7cm wide, one on each side of the spine between the rump and the shoulders. Only 3 maximum treatments per tick season annually. Make sure you re-treat when you first see ticks but not before at least six weeks after the previous treatment.

Breed Category

Pest Control

Buy Now AIRR

<https://www.airr.com.au/product/livestock/cattle/cattle-fly-lice/io-tick-off-pour-on/>

Safety Data Sheet

[Independents-Own-Tick-Off-Pour-On-Tick-Development-Inhibitor_valid-March-31](#)

Product Brochure

Missing Attachment

Product Label

[Tick-Off-5L---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery







Short Description

iO Tick Off Pour On is a tick development inhibitor and contains the powerful insect growth regulator (IGR), fluazuron 25g/L, which interferes with chitin synthesis of the larval stages of the cattle tick. Controls susceptible strains of cattle tick (*Boophilus microplus*) & Inhibits tick development within three days.

DrumMuster

1

Active Constituent

25g/L Fluazuron

Product Sizes

5L, 20L

Made in Australia

1

Pack size: Contains 5 x 500 g measure packs which it is illegal to sell separately.

Administration Instructions: This pack dissolves on contact with water. Open outer sachet. To dissolve in water, simply tip inner soluble pack directly into bucket and stir. Dispense as per mixing instructions advised on label.

Controls & Prevents:

ROUTINE TICK CONTROL (NSW, QLD, VIC, NT only): For routine control of specific types of ticks on cattle, apply HRC EXITRAZ at the following intervals:

CATTLE TICK (*Rhipicephalus* [*Boophilus*] *microplus*): intervals of 19 to 21 days.

BUSH TICK (NZ CATTLE TICK) (*Haemaphysalis longicornis*): intervals of 7 to 21 days.

PARALYSIS TICK (*Ixodes holocyclus*): intervals of 7 to 10 days. DO NOT leave longer intervals between treatments when ticks are active. For further information consult your Stock Inspector, District Veterinary Officer or Tick Extension Officer.



Breed Category

Pest Control

Buy Now AIRR

<https://www.airr.com.au/product/livestock/cattle/cattle-fly-lice/io-tick-off-pour-on/>

Safety Data Sheet

[Independents-Own-Tick-Off-Pour-On-Tick-Development-Inhibitor_valid-March-31](#)

Product Brochure

Missing Attachment

Product Label

[Tick-Off-5L---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery







Short Description

iO Tick Off Pour On is a tick development inhibitor and contains the powerful insect growth regulator (IGR), fluazuron 25g/L, which interferes with chitin synthesis of the larval stages of the cattle tick. Controls susceptible strains of cattle tick (*Boophilus microplus*) & Inhibits tick development within three days.

DrumMuster

1

Active Constituent

25g/L Fluazuron

Product Sizes

5L, 20L

Made in Australia

1

Nil meat and milk withholding provides flexibility particularly if clearing cattle of ticks prior to sale. Following treatment, cattle will be clean of ticks within 3-4 days, with larval ticks prevented from attaching for up to 7 days post treatment.

Active Constituent: 125 g/L Amitraz

Available in the following sizes: 400ML & 2L

Application: Coarse Spray with pump

Dosing & Administration

LIVESTOCK	PEST	STATE	RATE	CRITICAL COMMENTS
Cattle	Cattle Tick [Rhipicephalus (Boophilus) microplus]	NSW, QLD, WA and NT only	400mL per 200L water	Treat at intervals of 19-21 days Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick.Refer also to General Directions
	Paralysis Tick (Ixodes holocyclus)	NSW, QLD, WA and NT only	400mL per 200L water	Treat at intervals of 7-10 days Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick.Refer also to General Directions
	Bush Tick (NZ Cattle Tick) (Haemaphysalis longicornis)	NSW, QLD, WA, NT and VIC	400mL per 200L water	Treat at intervals of 7-21 days Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick.Refer also to General Directions
Sheep, goats, deer and certain circus animals	Cattle Tick [Rhipicephalus(Boophilus) microplus]	NSW only	400mL per 200L water	Treat animals on quarantined holdings at 14 to 21 day intervals. Treat animals being moved at 3 to 7 day intervals.
Pigs	Mange (Sarcoptes scabiei var. suis)	All States	40 mL per 10L water	Remove feed and bedding from pens, cover drinking bowls and clean out pen. Spray pigs with minimum 2L spray wash, especially inside ears and legs, under jowls and areas covered by scabs. Replace discarded bedding with clean material. Repeat treatment after 7-10 days. Refer also to General Directions.

Breed Category
Pest Control



Buy Now AIRR

<https://www.airr.com.au/product/livestock/cattle/cattle-fly-lice/io-tick-off-pour-on/>

Safety Data Sheet

[Independents-Own-Tick-Off-Pour-On-Tick-Development-Inhibitor_valid-March-31](#)

Product Brochure

Missing Attachment

Product Label

[Tick-Off-5L---PROOF-1](#)

Custom Product Download

Missing Attachment

Product Gallery







Short Description

iO Tick Off Pour On is a tick development inhibitor and contains the powerful insect growth regulator (IGR), fluazuron 25g/L, which interferes with chitin synthesis of the larval stages of the cattle tick. Controls susceptible strains of cattle tick (*Boophilus microplus*) & Inhibits tick development within three days.

DrumMuster

1

Active Constituent

25g/L Fluazuron

Product Sizes

5L, 20L

Made in Australia

1