

DIRECTIONS FOR USE

RESTRAINTS: DO NOT place baits in the open unless in bait stations.
DO NOT exceed 3 metres between baits for mice or 9 metres for rats.
DO NOT remove bait from paper sachet.
DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings	Mice (Mus musculus)	1-3 bait sachets per bait point depending on level of infestation	Place bait points wherever droppings, damage or other signs of rodent activity are seen, but not more than 2m away from buildings. For Mice: Place single baits at 2-3 m intervals in infested areas. For Rats: Place 1 to 3 baits at intervals of 5-9m in infested areas.
	Norway Rat (Rattus norvegicus)		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

The baits are ready to use – do not unwrap individual baits. Rats and mice will chew through the wrapper.

Note: One feed is sufficient to kill rats and mice. Ratshot Final Kill Paste can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations are used wherever baits are to be placed.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTIONS

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT feed the bait to wild or domestic birds. DO NOT apply directly on or near exposed food or food utensils, food preparation areas, livestock feed or water, or food storage areas.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



RATSHOT

FINAL KILL™ RODENTICIDE PASTE



ONE FEED KILLS

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM

READY TO USE BAIT • In a palatable sachet for the control of brown or norway rats (Rattus norvegicus) and Mice, including those resistant to other anti-coagulants.

FOR USE IN AND AROUND BUILDINGS

- ✓ Contains Denatonium Benzoate as a Bittering Agent
- ✓ One Feed Kills
- ✓ Ready to Use Edible Sachet



330 BAITS

CONTENTS

5

KG/NET

★★★★★

iO/iO

GUARANTEE

PROUDLY



OWNED



Freezone® In your Control.™

Contains (330 x 15g) bait sachets which it is illegal to sell separately

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT place baits in locations that are accessible to domestic animals, livestock, birds or other animals.
DO NOT use this product to control native animals. In areas where possums and other non-target animals frequent, baits should be secured inside proper bait stations or bait stations constructed from drainpipes or boxes.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT inhale dust. DO NOT touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Vitamin K1 (Phytomenadione) is antidotal.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet available from the supplier or visit www.freezone.net.au

Ratshot Final Kill Paste contains Denatonium benzoate (a human taste deterrent) to help prevent accidental human consumption.

NOTICE: Freezone Public Health Pty Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Freezone Public Health Pty Ltd, or under abnormal conditions.

Freezone, Ratshot & Final Kill are registered trademarks of Freezone Public Health Pty Ltd

APVMA Approval No.: 69210/60254

Batch Number

Date of Manufacture

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.



Freezone Public Health Pty Ltd
18 Gilpin Street, Shorncliffe, QLD 4017
Telephone: +61 (07) 3869 4436

