

265mm



177mm

PEEL BACK
LABEL TO REVEAL STRAP

General Directions:

iO Tick-Off Pour-On Treated Areas – Introduction of New Cattle: Tick infested cattle introduced onto properties where a iO Tick-Off Pour-On treatment regime is in place should be treated with iO Tick-Off Pour-On prior to release.

Handling of Treated Animals: Wear protective clothing recommended under Safety Directions, if handling treated cattle within 24 hours of application.

For local tick control recommendations contact your local department of agriculture.

Cleaning Instructions: After use wash iO Tick-Off Pour-On applicator with detergent and hot water. Dispose of the rinsates in a disposal pit away from desirable plants and their roots and watercourses.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.

MILK: DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI)
DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the registrant on 03 5820 8400 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS:
Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking and smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID:
If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

ADDITIONAL USER SAFETY INFORMATION:
Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION: DO NOT contaminate fish ponds, dams or rivers with the product or used containers. This product is very toxic to aquatic invertebrates.

DISPOSAL: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE: Store below 30°C (room temperature). Store in the tightly closed original container in a dry, cool place. Store out of direct sunlight.

APVMA APPROVAL NO.: 81471/121558

DIRECTIONS FOR USE:

Restraints: DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

Contraindications: This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.

Dosage and Administration:

ANIMALS & PEST	STATE	BODYWEIGHT (KG)	DOSE (ML)	NO. TREATED PER 5 L PACK
Beef cattle after weaning	QLD, NSW, WA & NT only	101-150	9	555
		151-200	12	416
Cattle Tick (<i>Rhipicephalus microplus</i>)		201-250	15	333
251-300		18	277	
301-350		21	238	
351-400		24	208	
401-450		27	185	
451-500		30	166	
501-550		33	151	
551-600		36	138	
601-650		39	128	
Over 650		3 mL per 50 kg		

CRITICAL COMMENTS

Retreat when you first see ticks but not before at least 6 weeks after the previous treatment.

Maximum number of treatments per annual tick season: 3.

Apply in two bands, about 7 cm wide, one on each side of the spine between the rump and the shoulders with an approved applicator. DO NOT use any other applicator.

The weight of cattle should be determined by either scales or a weigh-band. Dose individuals according to the heaviest animal in the mob. DO NOT under dose.

There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves.

iO Tick-Off Pour-On should not be applied to show cattle as a light scurfiness may be visible for some time after treatment.

iO Tick-Off Pour-On is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of iO Tick-Off Pour-On commences within 3 days of treatment.

Not suitable for cleaning cattle moving into 'gazetted' cattle tick areas.



NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

**Combustible liquid.
Causes skin irritation.
Causes serious eye damage.
May cause respiratory irritation.
May damage fertility or the unborn child.**

Avoid breathing fumes, mists, vapours or spray.

Use only outdoors or in a well-ventilated area.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

Store locked up.



B:

EXP:



179768-3

265mm

177mm