



BLADE

EQUINE WORMER

ACTIVE CONSTITUENTS: 100 mg/g PRAZIQUANTEL, 8 mg/g IVERMECTIN



**30
ML**



For the treatment and control of roundworms (including arterial larval stages of *Strongylus vulgaris* and benzimidazole resistant small strongyles), tapeworms, bots and skin lesions (summer sores) caused by *Habronema* and *Draschia spp.* and *Onchocerca spp.* microfilariae (cutaneous onchocerciasis).

At the recommended dose level of 1 mL/40 kg, iO BLADE EQUINE WORMER controls the following parasites:

Tapeworms: *Anoplocephala perfoliata*, *Anoplocephala magna*, *Paranoplocephala mammillana* (adult and immature heads and segments)

Large Strongyles: *Strongylus vulgaris* (adults and arterial larval stages), *Strongylus edentatus* (adults and tissue stages), *Strongylus equinus* (adults) and *Triodontophorus spp.* (adults)

Small Strongyles: Including benzimidazole resistant strains *Cyathostomum spp.*, *Cylicocyclus spp.*, *Cylicostephanus spp.*, *Cylicodontophorus spp.*, *Gyalocephalus spp.*

Pinworms: *Oxyuris equi* (adult and immature)

Ascarids: *Parascaris equorum* (adult and immature)

Hairworms: *Trichostrongylus axei* (adult)

Large Mouth Stomach Worms: *Habronema muscae* (adult)

Neck Threadworms: *Onchocerca spp.* (microfilariae)

Bots: *Gasterophilus spp.* (oral and gastric stages)

Lungworms: *Dictyocaulus arnfieldi* (adult and immature)

Intestinal Threadworms: *Strongyloides westeri* (adult)

iO BLADE EQUINE WORMER is also effective for the control of skin lesions caused by *Habronema* and *Draschia spp.* cutaneous larvae (summer sores) and *Onchocerca spp.* microfilariae (cutaneous onchocerciasis).

For SDS or technical information, please visit independentsown.com.au

