

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

MOXISTAR®

LONG ACTING INJECTION FOR CATTLE

ACTIVE CONSTITUENT: 100 MG/ML MOXIDECTIN 500 mL

For the treatment and control of moxidectin sensitive internal and external parasites in cattle.

Read the carton for full instructions.

DIRECTIONS FOR USE

Restraints:

DO NOT USE in cattle that may be exported live.

MILK: DO NOT USE in lactating cows or within 80 days of calving where milk may be used or processed for human consumption.

INJECT ONLY by subcutaneous injection into the back of the ear (as directed below). Do not inject anywhere else in the animal.

Injection sites anywhere else on the animal may result in injection site residues that exceed approved limits at the Withholding Period or Export Slaughter Interval.

Dosage and Administration:

Discard unused portion after 6 months of first broaching container.

Dose Rate: 1 mg moxidectin/kg liveweight (1 mL/100 kg liveweight)

Administer SC into the back of the ear towards its base. Avoid IV injection. DO NOT administer anywhere else on the animal. DO NOT use in cattle less than 100 kg or greater than 500 kg. Use a 5 mL injector gun fitted with a 16 or 18 gauge, 12 to 18 mm needle.

Re-treatment Interval: DO NOT retreat with MOXISTAR LONG ACTING INJECTION FOR CATTLE for at least 56 days after administration.

WITHHOLDING PERIODS

MEAT: 56 days.

MILK: 80 days.

TRADE ADVICE – EXPORT
SLAUGHTER INTERVAL (ESI)
108 days.

DO NOT USE in cattle that may be exported live.



THE HUNTER RIVER CO.

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the nose and throat. Will irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ENVIRONMENTAL PROTECTION

Moxidectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Dispose of container by wrapping with paper and putting in garbage. Discarded needles should immediately be placed in a designated and appropriately labelled "sharps" container.

STORAGE

Store below 30°C (room temperature). Protect from light.

BATCH:

EXP:

May cause respiratory irritation.

Causes damage to organs through prolonged or repeated exposure.

Do not get in eyes, on skin, or on clothing.

Wash contacted areas thoroughly after handling.

Do not eat, drink or smoke when using this product.

Wear protective gloves, protective clothing and eye or face protection.

Take off contaminated clothing and wash before reuse.

IF SWALLOWED, rinse mouth. DO NOT induce vomiting.

IF ON SKIN: Wash with plenty of soap and water.