

GENERAL INSTRUCTIONS

Most deaths occur 4 to 7 days after consumption of bait. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Record the number and locations of bait stations for purpose of replenishment and later removal.

Collect and dispose of all dead animals and rodent faeces, by burning or burial. When rodent activity has ceased, ensure uneaten bait is removed and disposed of as per disposal instructions.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

DO NOT contaminate dams, streams, rivers, waterways and drains with the bait or its used container. The bait and dead or affected rodents are hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock, birds or other non-target wildlife. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces e.g. drains. Regularly search for, remove and destroy dead or affected rodents by burning or burial. Ensure uneaten baits are removed and disposed of with empty containers when rodent activity has ceased.

STORAGE AND DISPOSAL:

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto

baiting site. Break, crush or puncture and bury empty container in an approved waste management facility. If an approved waste management facility is not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

PRECAUTION: KEEP OUT OF REACH OF CHILDREN

DO NOT place bait in areas where there is a possibility of contaminating food, livestock feed, water or surfaces that come into direct contact with food.

OTHER LIMITATIONS: THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES



31 631F-1

Freezone™ In your Control.™

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

RATSHOT^G
QUICKSHOTTM RODENTICIDE


io
Independents Own



ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)

READY TO USE GRAIN BAIT FOR USE IN COMMERCIAL, AGRICULTURAL AND DOMESTIC BUILDINGS

- ✓ Kills Rats and Mice
- ✓ Australian Cracked Wheat



CONTENTS

2

KG/NET

★★★★★

io
10

GUARANTEE

PROUDLY



OWNED & MADE



DIRECTIONS FOR USE RESTRAINTS:

DO NOT exceed 3m between bait stations for mice or 9m for rats.

DO NOT place bait stations in open.

DO NOT apply bait to crops.

BAITING STRATEGY:

Eliminate, as far as practicable, all other food sources. Bait the infested area. Place bait stations under cover along walls or rafters or in dark sheltered spots where there are signs of activity. Inspect bait stations daily. Replace eaten baits. If quickly eaten, increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers. **SITUATION:** In and around agricultural, commercial and domestic buildings (within 2 metres). Mice require several bait stations, 50g per station (0.2g will kill a mouse). Rats require larger quantities of bait in one or two locations, 100g - 200g per station in small populations (1.4g will kill a rat)*. For larger

populations more baiting points will be required. For effective control continue until feeding ceases.

*Based upon 25g mouse and 250g rat.

SAFETY DIRECTIONS:

Poisonous if swallowed. DO NOT inhale dust. DO NOT touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

SAFETY DATA SHEET:

Additional information is listed on the Safety Data Sheet available from the supplier or visit www.freezone.net.au

APVMA APPROVAL NO.:
84215/109902

io
Independents Own

Batch Number

Date of Manufacture

Freezone Public Health Pty Ltd
18 Gilpin Street, Shorncliffe, QLD 4017
Telephone: +61 (07) 3869 4436

Freezone

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.