

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY**



# **FLYK<sup>®</sup> EXTRA**

**SPRAY-ON SHEEP BLOWFLY TREATMENT**

**ACTIVE CONSTITUENT: 65 g/L DICYCLANIL**

For the protection for up to 29 weeks against strike by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*) when applied by spraying on sheep.

For the protection against strike by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*) during the wound healing process for sheep with mulesing and marking wounds.

**Made for AIRR**  
74-76 Drummond Rd  
Shepparton VIC 3630  
P 03 5820 8400  
independentsown.com.au

EXPIRY:

BATCH:

**CONTENTS**

**20**

**LITRES**




**iO / 10**

**GUARANTEE**

**PROUDLY**



**OWNED  
& MADE**



XXXXXXXX

iO FLYK EXTRA contains dicyclanil, a member of the pyrimidine group of chemicals. It is effective against strains of *Lucilia cuprina* that are sensitive to dicyclanil. It is not as effective against strains of *Lucilia cuprina* that develop resistance to dicyclanil. Resistance may develop to any chemical.

Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used.

#### DIRECTIONS FOR USE

##### Restrictions:

**DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT USE on sheep less than 4 weeks after shearing. DO NOT apply at the same time as any other backline product.**

**Re-treatment interval: DO NOT re-treat animals for a minimum of 9 weeks after last treatment.**

**Dosage and Administration: Shake well before use.**

**Use the contents within 12 months of first opening of the container. Discard the unused portion.**

#### SAFETY DIRECTIONS

May irritate eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product, wear elbow-length rubber gloves. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

#### ADDITIONAL USER SAFETY INFORMATION

Sheep Rehhandling Interval: After treatment wait until sheep are dry before rehhandling. Additional information is listed in the safety data sheet.

#### ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

#### DISPOSAL





Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.




#### STORAGE

Store below 30°C (room temperature).

Do not freeze. Protect from light. Store in original container tightly closed in a dry, cool place.

APVMA Approval No.: 91207/133579

FOR THE PREVENTION OF BODY STRIKE (DIAGRAM 1)				CRITICAL COMMENTS	
WEIGHT (kg)	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	Diagram 1.	
10-20	10	10	1	2 BANDS 	
21-30	12	12	1	4 BANDS 	
31-50	15	7.5	2		
>50	18	9	2		
FOR THE PREVENTION OF BREECH STRIKE (DIAGRAM 1)					
WEIGHT (kg)	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS		
10-20	10	10	1		
21-30	12	12	1		
31-50	15	7.5	2		
>50	18	9	2		
FOR THE PREVENTION OF POLL STRIKE (DIAGRAM 1)					
Rams and sheep susceptible to poll strike should be treated. Apply 5 mL of iO FLYK EXTRA to the areas requiring protection, around the poll and horns.					
FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (DIAGRAM 2)					
Apply two bands, one over each side of the mulesing wound and overlapping 25 mm onto the surrounding wool.					
WEIGHT (kg)	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS		
Up to 30	8	4	2		
31-50	10	5	2		
FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (DIAGRAM 3)					
<ul style="list-style-type: none"> <li>When tail has been amputated, apply 4 mL over the crutch ensuring sufficient coverage of the open wound.</li> <li>When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied.</li> </ul> <p>In case of ram lambs, a further 4 mL can be sprayed over and around the scrotum area after the rubber ring has been applied.</p>					
WEIGHT (kg)	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS		
Ram lambs	8	4	2		
Ewe lambs	4	4	1		



- Use only a spray-on applicator gun recommended by the manufacturer.
- Ensure the applicator gun is clean.
- Calibrate applicator gun using an accurate volume measure.
- Ensure all areas to be protected are treated.
- When preventing body strike ensure bands overlap along the midline (Diagram 1).
- When protecting mulesing wounds from fly strike, take special care to treat the wound and the surrounding 25 mm of wool (Diagram 2).
- When protecting marking wounds from fly strike, take special care to treat the wound and surrounding wool (Diagram 3).
- This product is not recommended for treating existing strikes. Use a product registered as a dressing for existing strikes.
- Heavy rain following the application of this product could diminish the period of protection.
- iO FLYK EXTRA contains a scourable colouring agent for easy band identification.
- Withholding periods must be observed following application of iO FLYK EXTRA for any purpose including after mulesing/markings as finite residues will occur.



**3082, ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S., PG III**

#### General Directions:

iO FLYK EXTRA provides season long protection to sheep against body, breech and poll strike for up to 29 weeks and may be used on sheep with any length wool from 4 weeks after shearing.

iO FLYK EXTRA also protects mulesing and marking wounds on lambs against fly strike during the wound healing process. The dose volume is determined by the weight of sheep being treated. Apply iO FLYK EXTRA according to the diagrams and tables in the Directions for Use. Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by the manufacturer. iO FLYK EXTRA may be applied to all breeds of sheep to obtain protection against fly strike. iO FLYK EXTRA is applied as a low-volume treatment onto areas where protection is required. When applying iO FLYK EXTRA, the nozzle of the applicator gun should be high enough above the fleece (approximately 15-25 cm), to achieve a 15 cm wide band at each pass for protection from body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool.

The product should not be applied during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for mulesing/markings to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis. Significant soiling with dags, mud, dust or urine may reduce efficacy. Where significant dags are present, a light crutching is recommended prior to treatment. Protection period may be reduced if sheep scour heavily following application.

Not to be mixed with any other product.  
Not to be diluted.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS**  
**MEAT: DO NOT USE less than 14 days before slaughter for human consumption.**

**MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.**

**WOOL: DO NOT USE less than 3 months before shearing or fibre collection.**

**TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI)**  
**DO NOT USE less than 9 weeks before slaughter for export. Before using this product, confirm the current ESI from the manufacturer on (03) 5820 8400 or the APVMA website (www.apvma.gov.au/residues).**

**Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.**

**Very toxic to aquatic life with long lasting effects.**  
Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling.  
**IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.**  
Collect spillage.  
In case of fire, use carbon dioxide, dry chemical, foam. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used.  
Store locked up and in a well-ventilated place. Keep cool.

XXXXXXXXXX