

ABASTAR

PLUS SELENIUM LV | ORAL DRENCH FOR SHEEP

ACTIVE CONSTITUENTS:

2.0 G/L Abamectin
1.0 G/L Selenium (as sodium selenate)

For the treatment and control of abamectin sensitive strains of gastrointestinal roundworms and lungworm in sheep, including benzimidazole and levamisole resistant strains.

For the treatment and control of nasal bot in sheep. For the treatment and prevention of selenium deficiencies in sheep.

TARGET PARASITES

Abastar Plus Selenium LV contains abamectin, a member of the avermectin family of drenches. It is effective against abamectin sensitive strains of the following internal parasites:

Adult and immature gastrointestinal roundworms:

Barber's pole worm - *Haemonchus contortus*
Small brown stomach worms - *Ostertagia spp.*
Black scour worms - *Trichostrongylus spp.*
Small intestinal worms - *Cooperia spp.*
Thin-necked intestinal worms - *Nematodirus spp.*
Large bowel worm - *Oesophagostomum venulosum*

Lungworm:

Dictyocaulus filaria

For the control of Sheep Nasal Bot - *Oestrus ovis*

For the treatment and prevention of selenium deficiencies in sheep.

RECOMMENDED APPLICATORS:

30 mL Variable Applicator

WITHHOLDING PERIODS:

Meat: DO NOT USE less than 14 days before slaughter for human consumption.

Milk: DO NOT USE in lactating ewes where milk may be used or processed for human consumption

Trade Advice - Export Slaughter Interval (ESI): An ESI has not been established for this product.

For SDS or technical information go to:
hunterriverco.com.au

DOSAGE & ADMINISTRATION:

1mL HRC Abastar Plus Selenium LV per 10kg bodyweight.
Drench sheep orally, using an appropriate drench gun.

DOSE RATE TABLE:

BODYWEIGHT	DOSE (ML)	NO. OF SHEEP TREATED
16 - 20	2.0	5000
21 - 25	2.5	4000
26 - 30	3.0	3333
31 - 35	3.5	2857
36 - 40	4.0	2500
41 - 45	4.5	2222
46 - 50	5.0	2000
51 - 55	5.5	1818
56 - 60	6.0	1666
61 - 65	6.5	1538
66 - 70	7.0	1428
71 - 75	7.5	1333
Animals heavier than 75 kg to be dosed at 1 mL per 10 kg.		

