

FRONT

100mm



The packaging design for Levistar LV Oral Drench features a large, stylized letter 'L' in a blue-to-green gradient. A teal silhouette of a horse is positioned behind the 'L'. To the left of the 'L', the text '10 L' is written in a bold, blue font. A dotted line indicates a fold line for the top of the 'L'. On the right side, the word 'Levistar' is written in a large, blue, italicized font, with 'LV' in a smaller, blue font. Below this, 'ORAL DRENCH' is written in a bold, green font. The text 'ACTIVE CONSTITUENT: 80 g/L LEVAMISOLE HYDROCHLORIDE (equivalent to 68 g/L LEVAMISOLE)' is in a smaller, black font. Below that, the text 'For the control of levamisole sensitive Round Worms including Lungworms in Sheep and Cattle.' is in a green font. At the bottom right, there are two icons: an orange horse head and a teal sheep head. To the right of these icons is the Hunter River Co. logo, which consists of a stylized green and blue wave. The text '23559F-2' is printed vertically on the right side of the packaging.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

**Levistar LV**  
**ORAL DRENCH**

ACTIVE CONSTITUENT:  
80 g/L LEVAMISOLE HYDROCHLORIDE (equivalent to 68 g/L LEVAMISOLE)

For the control of levamisole sensitive Round Worms  
including Lungworms in Sheep and Cattle.

10 L

HUNTER RIVER CO.

23559F-2

200mm

# LEFT SIDE

For the control of levamisole sensitive gastrointestinal roundworms and lungworms in sheep and cattle. LEVISTAR LV contains levamisole, a member of the imidazothiazole (clear) group of anthelmintics (drenches) and is effective in cattle and sheep against: mature and immature Barber's Pole Worm (*Haemonchus* spp.), Stomach Hair Worm (*Trichostrongylus axei*), Black Scour Worm (*Trichostrongylus* spp.), Small Intestinal worm (*Cooperia* spp.), Thin Necked Intestinal Worm (*Nematodirus* spp.), Hookworm (*Bunostomum* spp.), Large Mouthed Bowel Worm (*Chabertia ovina*), Strongyloides, Nodule and Large Bowel Worms (*Oesophagostomum* spp.), Large Lungworm (*Dictyocaulus* spp.) and Small Brown Stomach Worm (*Teladorsagia* (*Ostertagia* spp.) in sheep, and mature (adult only) *Ostertagia* spp. in cattle under 12 months of age. Where *Ostertagia* are the problem in cattle over 12 months of age, STARMEC POUR-ON FOR BEEF AND DAIRY CATTLE is the recommended product.

Controls both susceptible and benzimidazole resistant strains of Barber's pole worm (*Haemonchus* species) and Black scour worm (*Trichostrongylus* species) in sheep.

Resistance may develop to any chemical.

### DIRECTIONS FOR USE

#### Contraindications:

This product is contraindicated for use in dogs and horses.

#### Precautions:

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress.

It is desirable to hold pregnant cows and heifers in the yards or a small paddock for 2-3 hours after treatment.

Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching.

Recommended dose should not be exceeded, except under veterinary supervision.

#### Side Effects:

Cattle may show temporary signs of stimulation and friskiness when released from the yards soon after treatment. Other slight side effects of brief duration such as salivation and shivering may be seen.

**Dosage and Administration:**  
LEVISTAR LV is given as a drench by mouth in cattle and sheep.

**Dose rates for CATTLE:** 5 mL per 50 kg liveweight. The liveweight should be estimated carefully so that the dose can be calculated accurately to ensure optimum efficiency with economy. The following table may be used as a guide.

Cattle heavier than 650 kg should be dosed at 5 mL per 50 kg.

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

**Dose rates for SHEEP:** 1 mL per 10 kg liveweight. Always dose on a liveweight basis. Check the accuracy of drenching equipment before and during drenching procedures. The following table may be used as a guide.

Sheep heavier than 80 kg should be dosed at 1 mL per 10 kg.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate to avoid excessive overdosing.

23559LEFT-2

LIVEWEIGHT (KG)	DOSE (ML)	NO. OF CATTLE TREATED
Up to 50	5	2,000
51 – 100	10	1,000
101 – 150	15	666
151 – 200	20	500
201 – 250	25	400
251 – 300	30	333
301 – 350	35	285
351 – 400	40	250
401 – 450	45	222
451 – 500	50	200
501 – 550	55	181
551 – 600	60	166
601 – 650	65	153

LIVEWEIGHT (KG)	DOSE (ML)	NO. OF SHEEP TREATED
Up to 10	1	10,000
11 – 20	2	5,000
21 – 30	3	3,333
31 – 40	4	2,500
41 – 50	5	2,000
51 – 60	6	1,666
61 – 70	7	1,428
71 – 80	8	1,250

# RIGHT SIDE

#### General Directions:

If drench (anthelmintic) failure with this preparation is suspected, seek veterinary confirmation and notify the manufacturer. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any parasite treatment is used.

#### WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 3 days before slaughter for human consumption.**

**MILK: Zero (0) days.**

#### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export. Before using this product, confirm the current ESI from The Hunter River Company Pty Limited on (03) 5820 8400 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**

#### SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wear rubber gloves.

#### FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

#### ADDITIONAL USER SAFETY

Additional information is listed in the Safety Data Sheet.

#### ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers or streams with the drench or used container.

#### DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately at the drumMUSTER

collection site. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### STORAGE

Store below 30°C (room temperature) in the closed, original container in a well-ventilated area, as cool as possible. Protect from light.

#### Harmful in contact with skin.

Wash contacted areas thoroughly after handling.

Do not eat, drink or smoke when using this product.

Wear protective gloves, protective clothing and eye or face protection.

Wash contaminated clothing before reuse.

**IF SWALLOWED:** Immediately call a POISON CENTRE or doctor.

Rinse mouth. Do NOT induce vomiting.

**IF ON SKIN:** Wash with plenty of soap and water.

Not combustible. Use extinguishable media suited to burning materials.

BATCH:

APVMA Approval No. 86346/138226

EXP:



23559RIGHT-2

100mm

The Hunter River Company Pty Limited | ACN 133 798 615 | 74-76 Drummond Road, Shepparton VIC 3630 | Ph: (03) 5820 8400

150mm