

EXITRAZ EC

Active Constituent: 125 g/L Amitraz
Also contains: 600 g/L Hydrocarbon liquid (solvent)

EASY TO USE LIQUID CONCENTRATE FOR TICK CONTROL IN BEEF AND DAIRY CATTLE



FOR THE CONTROL OF

- Organophosphate and Synthetic Pyrethroid resistant Cattle Tick
- Paralysis Tick and Bush Tick on Beef and Dairy Cattle, and
- Cattle Tick on Sheep, Goats, Deer and Certain Circus animals in NSW only, and
- Mange on Pigs.

APPLICATOR

Coarse Spray – with pump

WITHHOLDING PERIODS

MEAT: Zero (0) days MILK: Zero (0) days ESI: Zero (0) days

STORAGE

Store below 30°C (Room Temperature) in the closed original container in a dry, wellventilated area, out of direct sunlight.

COMPARABLE PRODUCT

Coopers® Amitik® EC.

DOSING & ADMINISTRATION

LIVESTOCK	PEST	STATE	RATE	CRITICAL COMMENTS	
Cattle	Cattle Tick [Rhipicephalus (Boophilus) microplus]	NSW, QLD, WA and NT only	400mL per 200L water	Treat at intervals of 19-21 days	Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick. Refer also to General Directions
	Paralysis Tick (lxodes holocyclus)			Treat at intervals of 7-10 days	
	Bush Tick (NZ Cattle Tick) (Haemaphysalis longicornis)	NSW, QLD, WA, NT and VIC		Treat at intervals of 7-21 days	
Sheep, goats, deer and certain circus animals	Cattle Tick [Rhipicephalus (Boophilus) microplus]	NSW only	400mL per 200L water	Treat animals on quarantined holdings at 14 to 21 day intervals. Treat animals being moved at 3 to 7 day intervals.	
Pigs	Mange (Sarcoptes scabiei var. suis)	All States	40 mL per 10L water	Remove feed and bedding from pens, cover drinking bowls and clean out pen. Spray pigs with minimum 2L spray wash, especially inside ears and legs, under jowls and areas covered by scabs. Replace discarded bedding with clean material. Repeat treatment after 7-10 days. Refer also to General Directions.	

AVAILABLE SIZES

400ML, 2L

For more information please visit hunterriverco.com.au

Coopers and Amitik are registered trade marks of Intervet Inc. a Delaware corporation.





