

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



fenbender 100[®]

ORAL DRENCH FOR CATTLE & HORSES

ACTIVE CONSTITUENT: 100 g/L FENBENDAZOLE

For the control of benzimidazole-sensitive mature and immature Roundworms (including inhibited *Ostertagia*), Lungworms and as an aid in the control of Tapeworms in Cattle. For the treatment and control of Roundworms, Bloodworms, Redworms and Pinworms in horses.

io FENBENDER 100[®] ORAL DRENCH FOR CATTLE AND HORSES contains Fenbendazole, a member of the benzimidazole family of chemicals.

It is effective against sensitive strains of the following internal parasites:

CATTLE: For the control of Benzimidazole-Sensitive mature and immature Roundworms (including inhibited *Ostertagia*), Lungworms and as an aid in the control of Tapeworms in cattle:

(*Haemonchus placei*) Barber's pole worm;
(*Trichostrongylus axei*) Stomach hair worm;
(*Trichostrongylus* spp) Intestinal hair worm or Black scour worm; (*Bunostomum phlebotomum*) Hookworm; (*Cooperia* spp) Small intestinal worm; (*Nematodirus* spp) Thin necked intestinal worm; (*Oesophagostomum radiatum*) Nodule worm; (*Dictyocaulus viviparus*) Lungworm, (*Ostertagia ostertagi*) Small brown stomach worm including 80% - 90% activity against Hypobiotic larvae (*Moniezia* spp) Tapeworm.

HORSES: For treatment and control of mature and immature Roundworms, Bloodworms, Redworms and Pinworms in horses:

(*Strongylus* spp) Large strongyle, (*Cyathostomum* spp, *Triodontophorus* spp.) Small strongyle, (*Parascaris equorum*) Ascarids, (*Oxyuris equi*) Pinworms.

Resistance may develop to any chemical.

DIRECTIONS FOR USE:**Restraints: NOT FOR USE IN GOATS****Dosage and Administration:****SHAKE WELL BEFORE USE.**

This product can be administered concurrently with vaccines, pour-on lice treatments and dipping operations.

CATTLE: 7.5 mg/kg Fenbendazole or 3 mL per 40 kg bodyweight.

Cattle in excess of 680 kg to be dosed at 3 mL per 40 kg bodyweight.

Bodyweight (kg)	Dose (mL)	No. of Animals Treated Per Pack		
		1 L	5 L	10 L
Up to 80	6	167	833	1666
81 - 120	9	133	556	1111
121 - 160	12	83	417	833
161 - 200	15	67	333	666
201 - 240	18	55	278	555
241 - 280	21	47	238	476
281 - 320	24	41	208	416
321 - 360	27	37	185	370
361 - 400	30	33	167	333
401 - 440	33	30	152	303
441 - 480	36	27	139	277
481 - 520	39	25	128	256
521 - 560	42	23	119	238
561 - 600	45	22	111	222
601 - 640	48	21	104	208
641 - 680	51	19	98	196

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Check accuracy of drenching gun before and during use. Rinse drenching guns after use.

HORSES: 10 mg/kg

Fenbendazole or 5 mL per 50 kg body weight. For the control of *Strongyloides westeri* in foals: 50 mg/kg or 5 mL/10 kg body weight. iO Fenbender 100 Oral Drench may be administered to horses by nasal tube, drenching gun or dosage syringe.

Worm parasites may develop resistance to Fenbendazole. If this is suspected, please contact your local veterinary practitioner or advise the manufacturer.

WITHHOLDING PERIODS:

CATTLE: MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: Zero (0) days.

HORSES: DO NOT USE less than 28 days before slaughter for human consumption.

TRADE ADVICE**EXPORT SLAUGHTER INTERVAL (ESI):**

CATTLE: DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of approval. Before using this product confirm the ESI from the distributor on (03) 5820 8400 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS: Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

FIRST AID INSTRUCTIONS: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY: Additional information is listed in the safety data sheet (SDS).

DISPOSAL (1L): Dispose of empty container by wrapping with paper and putting in garbage.

DISPOSAL (5L & 10L): Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE: Store below 30°C (Room temperature) with top secure - Avoid Freezing.

MADE FOR AIRR

74-76 Drummond Rd,
Shepparton VIC 3630

P. 03 5820 8400
independentsown.com.au

APVMA Approval Number: 81050/102239