

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



mectec plus

(IVERMECTIN PLUS CLORSULON)

BROAD-SPECTRUM ANTIPARASITIC INJECTION FOR CATTLE

ACTIVE CONSTITUENTS: 100 mg/mL CLORSULON, 10 mg/mL IVERMECTIN

For the treatment and control of ivermectin and clorsulon sensitive strains of internal and external parasites of cattle, including adult liver flukes.

500 mL

Read the carton for full instructions.

DIRECTIONS FOR USE

Restraints:

Re-treatment interval: DO NOT re-treat animals for 28 days after last treatment.

Dosage and Administration:

To be given by subcutaneous injection at the recommended dose of 1 mL per 50 kg liveweight (equivalent to 0.2 mg ivermectin per kg and 2 mg clorsulon per kg liveweight). Refer to carton for dosage table.

Inject under loose skin high on the neck, behind the ear in an area away from the more valuable carcass muscle. Loose skin on the neck in front of the shoulder is also a suitable site. This product is not to be used intravenously or intramuscularly.

WITHHOLDING PERIODS

MEAT: 28 days.

MILK: Zero (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): 42 days.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes, skin and clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ENVIRONMENTAL PROTECTION

Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Dispose of container and outer packaging by wrapping with paper and putting in garbage.

STORAGE

Store below 30°C (room temperature). Store bottle in carton to protect from light.

APVMA Approval No.: 90419/131992

BATCH:

EXPIRY:

MADE FOR AIRR 74-76 Drummond Rd, Shepparton, VIC 3630

P 03 5820 8400 | independentsown.com.au

23639C-5