

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



tripletec LV

3-WAY MINERALISED COMBINATION DRENCH FOR SHEEP AND CATTLE

ACTIVE CONSTITUENTS: 80 g/L LEVAMISOLE HYDROCHLORIDE, 45.3 g/L OXFENDAZOLE, 5 g/L COBALT
(as DISODIUM COBALT EDTA), 2 g/L ABAMECTIN, 1 g/L SELENIUM (as SODIUM SELENATE)

MADE FOR AIRR 74-76 Drummond Rd, Shepparton, VIC 3630 | P 03 5820 8400 | independentsown.com.au



For the treatment and control of internal parasites of sheep including those resistant to Macrocytic Lactone (ML), Benzimidazole (BZ) or Levamisole (LEV) drench families.

For the treatment and control of internal parasites of cattle including those resistant to Macrocytic Lactone (ML) drench family.

To supplement diets which may be deficient in selenium and cobalt.



10L

DIRECTIONS FOR USE

Restraints:

DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption. Re-treatment interval: DO NOT RE-TREAT animals for 28 days after last treatment.

Contraindications:

This product is contraindicated for use in lambs under 6 weeks of age or weighing 15 kg or less.

This product is contraindicated for use in calves under 16 weeks of age or weighing 100 kg or less.

This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) or if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian. This product should not be used in dogs, horses or any other pets.

This product is not recommended for use in goats as safety and efficacy have not been evaluated.

Precautions:

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching. Recommended dose should not be exceeded.

Use with care as selenium can have toxic effects, particularly in cattle and sheep with liver damage, including cattle and sheep that graze in areas where plants containing pyrrolizidine alkaloids are prevalent.

Safety in breeding cattle and sheep of either sex in pregnant or lactating cows or ewes has not been established.

Dosage and Administration:

SHAKE WELL BEFORE USE.

Before commencing drenching read the label carefully.

iO Tripletec LV should be administered orally using accurate drenching equipment. Check the accuracy of drenching equipment before and during drenching procedures. Clean equipment thoroughly before and after use.

Dose Rate: 1 mL per 10 kg bodyweight by a single oral administration.

SHEEP

Liveweight (kg)	Dose (mL)	Liveweight (kg)	Dose (mL)
15 or less	Do not treat	51 - 60	6
16 - 20	2	61 - 70	7
21 - 30	3	71 - 80	8
31 - 40	4	Over 80	1 mL / 10 kg
41 - 50	5		

Sheep heavier than 80 kg to be dosed at 1 mL per 10 kg. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into 2 or more lines may be appropriate, to avoid excessive overdosing.

CATTLE

Liveweight (kg)	Dose (mL)	Liveweight (kg)	Dose (mL)
100 or less	Do not treat	401 - 450	45
101 - 150	15	451 - 500	50
151 - 200	20	501 - 550	55
201 - 250	25	551 - 600	60
251 - 300	30	601 - 650	65
301 - 350	35	Over 650	1 mL / 10 kg
351 - 400	40		

Cattle heavier than 650 kg should be dosed at 1 mL per 10 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not underdose. Where there is a large variation in size within the group, draft into 2 or more lines based on body weight, to avoid excessive overdosing.

General Directions:

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: Cattle - DO NOT USE in lactating cows or within 28 days of calving where milk or milk products will be used for human consumption. If cows calve earlier than 28 days after treatment, milk may contain residues. This milk MUST NOT BE USED for human consumption, or supplied for processing for at least 28 days following treatment.

Calves fed this milk should not be slaughtered for human consumption within 21 days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): Cattle: DO NOT USE less than 21 days before slaughter for export.

Sheep: DO NOT USE less than 28 days before slaughter for export. Before using this product, confirm the current ESI from the distributor on (03) 5820 8400 or the APVMA website (apvma.gov.au/residues).